

## **PREVENTIVE CARE SERVICES**

#### Guideline Number: CDG.016.17

#### Effective Date: October 1, 2017

Table of Contents	
INSTRUCTIONS FOR USE	1
BENEFIT CONSIDERATIONS	1
COVERAGE RATIONALE	2
DEFINITIONS	5
APPLICABLE CODES	6
REFERENCES	45
GUIDELINE HISTORY/REVISION INFORMATION	46

Related Commercial Policies

- <u>Breast Imaging for Screening and Diagnosing</u> <u>Cancer</u>
- <u>Cardiovascular Disease Risk Tests</u>
- <u>Computed Tomographic Colonography</u>
- <u>Cytological Examination of Breast Fluids for Cancer</u> <u>Screening</u>
- <u>Genetic Testing for Hereditary Breast and/or</u> <u>Ovarian Cancer Syndrome (HBOC)</u>
- Preventive Medicine and Screening Policy
- Vaccines

## INSTRUCTIONS FOR USE

This Coverage Determination Guideline provides assistance in interpreting UnitedHealthcare benefit plans. When deciding coverage, the member specific benefit plan document must be referenced. The terms of the member specific benefit plan document [e.g., Certificate of Coverage (COC), Schedule of Benefits (SOB), and/or Summary Plan Description (SPD)] may differ greatly from the standard benefit plan upon which this Coverage Determination Guideline is based. In the event of a conflict, the member specific benefit plan document supersedes this Coverage Determination Guideline. All reviewers must first identify member eligibility, any federal or state regulatory requirements, and the member specific benefit plan coverage prior to use of this Coverage Determination Guideline. Other Policies and Coverage Determination Guidelines may apply. UnitedHealthcare reserves the right, in its sole discretion, to modify its Policies and Guidelines as necessary. This Coverage Determination Guideline is provided for informational purposes. It does not constitute medical advice.

UnitedHealthcare may also use tools developed by third parties, such as the MCG<sup>™</sup> Care Guidelines, to assist us in administering health benefits. The MCG<sup>™</sup> Care Guidelines are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.

#### BENEFIT CONSIDERATIONS

Before using this guideline, please check the member specific benefit plan document and any federal or state mandates, if applicable.

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HHS: Health and Human Services
- HRSA: Health Resources and Services Administration

For self-funded plans with SPD language other than fully-insured Generic COC language, please refer to the member specific benefit plan document for coverage.

#### Essential Health Benefits for Individual and Small Group

For plan years beginning on or after January 1, 2014, the Affordable Care Act of 2010 (ACA) requires fully insured non-grandfathered individual and small group plans (inside and outside of Exchanges) to provide coverage for ten categories of Essential Health Benefits ("EHBs"). Large group plans (both self-funded and fully insured), and small group ASO plans, are not subject to the requirement to offer coverage for EHBs. However, if such plans choose to provide coverage for benefits which are deemed EHBs, the ACA requires all dollar limits on those benefits to be removed on all Grandfathered and Non-Grandfathered plans. The determination of which benefits constitute EHBs is made on a state by state basis. As such, when using this guideline, it is important to refer to the member specific benefit plan document to determine benefit coverage.

#### COVERAGE RATIONALE

#### **Indications for Coverage**

#### Introduction

UnitedHealthcare covers certain medical services under the Preventive Care Services benefit. Effective for plan years on or after September 23, 2010, the federal Patient Protection and Affordable Care Act (PPACA) requires nongrandfathered health plans to cover certain "recommended preventive services" identified by PPACA under the Preventive Care Services benefit, without cost sharing to members when provided by Network physicians.

#### For Plan Years that begin on or after September 23, 2010

For non-grandfathered health plans, UnitedHealthcare will cover the recommended preventive services under the Preventive Care Services benefit as mandated by PPACA, with no cost sharing when provided by a Network provider. These services are described in the United States Preventive Services Task Force A and B recommendations, the Advisory Committee on Immunization Practices (ACIP) of the CDC, and Health Resources and Services Administration (HRSA) Guidelines including the American Academy of Pediatrics *Bright Futures* periodicity guidelines.

#### For Plan Years that begin on or after August 1, 2012

For non-grandfathered plans, UnitedHealthcare will cover for women the additional preventive care and screenings as required by the HHS Health Plan Coverage Guidelines for Women's Preventive Services for plan years that begin on or after August 1, 2012.

In addition to these mandated services, under the Preventive Care Services benefit, UnitedHealthcare also covers screening using CT colonography; prostate specific antigen (PSA); and screening mammography for adult women without age limits.

#### **Grandfathering for Preventive Care Services**

Grandfathered plans, as that term is defined under PPACA, are not required by law to provide coverage without cost sharing for preventive services; although a grandfathered plan may amend its plan document to voluntarily comply with the preventive benefit requirements under PPACA.

Grandfathered health plans will continue the benefits for preventive care that existed in the plan prior to September 23, 2010, without conforming to the federal mandate under PPACA, unless amended to comply with the federal requirements. Except where there are State mandates, a grandfathered plan might include member cost sharing or exclude some of the preventive care services identified under PPACA. Please refer to the member specific plan document for details.

Non-grandfathered plans are required to cover the preventive care services as defined in the PPACA at no cost sharing. Please refer to the member specific plan document for details.

## **Cost Sharing for Non-Grandfathered Health Plans**

**Network** Preventive Care Services that are identified by PPACA are required to be covered under the Preventive Care Services benefit with no member cost sharing (ie. covered at 100% of Eligible Expenses without deductible, coinsurance or copayment). Depending on the plan, Eligible Expenses for services from non-network providers may not equal the provider's billed charges (refer to plan's schedule of benefits).

**Note**: For Network providers, UnitedHealthcare has made a decision to also cover the "Additional Preventive Care Services" identified below with no member cost sharing.

**Non-Network** preventive care services are not part of the PPACA requirements. Many plans do not cover nonnetwork preventive care services. If a plan covers non-network preventive care services, the benefit for non-network is allowed to have member cost sharing. Please refer to the member specific plan document for non-network information.

#### Summary of Preventive Care Services Benefit

The following is a high-level summary of the services covered under the Preventive Care Services benefit (broken down by age/gender groups):

**All members**: Age- and gender-appropriate Preventive Medicine visits (Wellness Visits); all routine immunizations recommended by the Advisory Committee on Immunization Practices of the CDC.

**All members at an appropriate age and/or risk status:** Counseling and/or screening for: colorectal cancer; elevated cholesterol and lipids; certain sexually transmitted diseases; HIV; depression; high blood pressure; diabetes. Screening and counseling for alcohol abuse in a primary care setting; tobacco use; obesity; diet and nutrition.

## Women's Health

1. *Plan Years that Begin on or after September 23, 2010*: Screening mammography; cervical cancer screening including Pap smears; genetic counseling and evaluation for the BRCA breast cancer gene test; BRCA lab screening\* (effective October 1, 2013); counseling for chemoprevention for women at high risk for breast cancer; screening for gonorrhea, chlamydia, syphilis in defined high risk groups; osteoporosis screening. Screening pregnant women for bacteriuria; hepatitis B virus; Rh incompatibility; and instructions to promote and aid with breast feeding.

\*Prior Authorization for BRCA Testing:

- For most benefit plans, prior authorization requirements apply to BRCA lab screening.
- For medical necessity benefit plans: genetic counseling from an Independent Genetics Provider (see definition section) is required before UnitedHealthcare will approve prior authorization requests (effective January 1, 2016).
- 2. Plan Years that Begin on or after August 1, 2012: Preventive visits to include preconception and prenatal services; FDA-approved contraception methods and contraceptive counseling; human papillomavirus (HPV) DNA testing for women 30 years and older; breastfeeding support and counseling, and costs of breastfeeding equipment; domestic violence screening and counseling; annual human immunodeficiency virus (HIV) screening and counseling; annual sexually-transmitted infection counseling; and screening for gestational diabetes for all pregnant women that have no prior history of diabetes.

## Men's Health

Screening for prostate cancer for men age 40 and older; screening for abdominal aortic aneurysm in men 65-75 years old (USPSTF recommends this for males 65-75 years old who have smoked).

## **Pediatrics**

Screening newborns for hearing problems, thyroid disease, phenylketonuria, sickle cell anemia, and standard metabolic screening panel for inherited enzyme deficiency diseases. For children (at the appropriate age): Application of fluoride by a primary care provider, for prevention of dental cavities; screening for major depressive disorders; vision; lead; tuberculosis; developmental/autism; counseling for obesity.

## **Additional Preventive Care Services**

The following preventive care services are not currently required by PPACA. However, these services are covered under UnitedHealthcare's Preventive Care Services benefit.

- 1. Mammography (film and digital) screening for all adult women
- 2. Computed Tomographic Colonography (Virtual Colonoscopy) for screening for colon cancer
- 3. Osteoporosis Screening for all women (regardless of risk)
- 4. Prostate Cancer Screening for all men age 40+
- 5. Wellness / Physical Examinations for Adults (Age- and gender-appropriate)\*

\* See the Expanded Women's Preventive Health coding table below regarding specific services that are covered as well woman visits under PPACA, for plan years that begin on or after August 1, 2012.

## Preventive vs. Diagnostic Services

Certain services can be done for preventive or diagnostic reasons. When a service is performed for the purpose of preventive screening and is appropriately reported, it will be adjudicated under the Preventive Care Services benefit.

Preventive services are those performed on a person who:

- 1. has not had the preventive screening done before and does not have symptoms or other abnormal studies suggesting abnormalities; or
- 2. has had screening done within the recommended interval with the findings considered normal; or
- 3. has had diagnostic services results that were normal after which the physician recommendation would be for future preventive screening studies using the preventive services intervals.
- 4. has a preventive service done that results in a therapeutic service done at the same encounter and as an integral part of the preventive service (e.g., polyp removal during a preventive colonoscopy), the therapeutic service would still be considered a preventive service.

Examples include, but are not limited to:

- A woman had an abnormal finding on a preventive screening mammography and the follow up study was found to be normal, and the patient was returned to normal mammography screening protocol, then future mammography would be considered preventive.
- If a polyp is encountered during preventive screening colonoscopy, the colonoscopy, removal of the polyp, and associated facility, lab and anesthesia fees done at the same encounter are covered under the Preventive Care Services benefit.

When a service is done for diagnostic purposes it will be adjudicated under the applicable non-preventive medical benefit.

Diagnostic services are done on a person who:

- 1. had abnormalities found on previous preventive or diagnostic studies that require further diagnostic studies; or
- 2. had abnormalities found on previous preventive or diagnostic studies that would recommend a repeat of the same studies within shortened time intervals from the recommended preventive screening time intervals; or
- 3. had a symptom(s) that required further diagnosis; or
- 4. does not fall within the applicable population for a recommendation or guideline (e.g., someone who has a colorectal cancer screening due to a family history).

Examples include, but are not limited to:

- A patient had a polyp found and removed at a prior preventive screening colonoscopy. All future colonoscopies are considered diagnostic because the time intervals between future colonoscopies would be shortened.
- A patient had an elevated cholesterol on prior preventive screening. Once the diagnosis has been made, further testing is considered diagnostic rather than preventive. This is true whether or not the patient is receiving pharmacotherapy.
- If a Preventive service results in a therapeutic service at a later point in time, the Preventive Service would be adjudicated under the Preventive Care Services benefit and the therapeutic service would be adjudicated under the applicable non-preventive medical benefit.

## **Related Services**

Services that are directly related to the performance of a preventive service are adjudicated under the Preventive Care Services benefit. Examples include:

- All services for a preventive colonoscopy (e.g., associated facility, anesthesia, pathologist, and physician fees). The preventive benefit does **not** include a post-operative examination. Effective January 1, 2016, the preventive benefit includes a pre-operative examination / consultation prior to a preventive colonoscopy.
- Women's outpatient sterilization procedures (e.g., associated implantable devices, facility fee, as well as anesthesia, pathology, and physician fees) are considered to be related services and covered under the preventive benefit. Note the following:
  - a. The preventive benefit does **not** include a pre- or post-operative examination.
  - b. If a woman is admitted to an inpatient facility for another reason, and has a sterilization performed during that admission, the sterilization surgical fees (surgical fee, device fee, anesthesia, pathologist and physician fees), are covered under the preventive benefit. However, the facility fees are not covered under preventive benefits since the sterilization is incidental to and is not the primary reason for the admission.
  - c. For hysteroscopic fallopian tube occlusion sterilization procedures, the preventive benefit includes an outpatient, followup hysterosalpingogram to confirm that the fallopian tubes are completely blocked.
- 3. Blood drawing (venipuncture or finger or heel stick) is considered as payable under the preventive benefit if billed for a preventive lab service that requires a blood draw.

**Note**: However, that benefit adjudication is contingent upon accurate claims submission by the provider, including diagnosis, procedure, age and gender.

## **Covered Breastfeeding Equipment**

Personal-use electric breast pump:

- The purchase of a personal-use electric breast pump (HCPCS code E0603).
  - This benefit is limited to one pump per birth. In the case of a birth resulting in multiple infants, only one breast pump is covered.
  - A breast pump purchase includes the necessary supplies for the pump to operate.
- Replacement breast pump supplies necessary for the personal-use electric breast pump to operate. This includes: standard power adaptor, tubing adaptors, tubing, locking rings, bottles specific to breast pump operation, caps for bottles that are specific to the breast pump, valves, filters, and breast shield and/or splash protector for use with the breast pump.

Note: See Coverage Limitations and Exclusions section for non-covered items.

## Additional Information

- A new immunization that is pending ACIP recommendations, but is a combination of previously approved individual components, is eligible for adjudication under the preventive care benefit.
- Refer to the reimbursement policy titled <u>Preventive Medicine and Screening Policy</u> for situations which may affect reimbursement of preventive care services.
- The list of recommended preventive services covered will be updated as new recommendations and guidelines are issued, or as existing ones are revised or removed by the USPSTF, ACIP and the HRSA. Updates will occur no less frequently than required by PPACA.

## **Coverage Limitations and Exclusions**

- 1. Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- 2. Generally, the cost of drugs, medications, vitamins, supplements, or over the counter items are not eligible as a preventive care benefit. However, certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA, may be covered under the preventive benefit. For details, please refer to the member-specific pharmacy plan administrator.
- An immunization is not covered if it does not meet company Vaccine Policy requirements for FDA labeling (including age and/or gender limitations) and if it does not have definitive ACIP recommendations published in the CDC's Morbidity and Mortality Weekly Report (MMWR).
- 4. Examinations, screenings, testing, or immunizations are not covered when:
  - a. required solely for the purposes of career, education, sports or camp, travel (including travel immunizations), employment, insurance, marriage or adoption, or
  - b. related to judicial or administrative proceedings or orders, or
  - c. conducted for purposes of medical research, or
  - d. required to obtain or maintain a license of any type.
- 5. Services that are investigational, experimental, unproven or not medically necessary are not covered. Please see applicable Medical Policies for details.
- 6. Breastfeeding equipment and supplies not listed in the Indications for Coverage section above. This includes, but is not limited to:
  - a. Manual breast pumps and all related equipment and supplies.
  - b. Hospital-grade breast pumps and all related equipment and supplies.
  - c. Equipment and supplies not listed in the <u>Covered Breastfeeding Equipment</u> section above, including but not limited to:
    - o Batteries, battery-powered adaptors, and battery packs.
    - o Electrical power adapters for travel.
    - o Bottles which are not specific to breast pump operation. This includes the associated bottle nipples, caps and lids.
    - o Travel bags, and other similar travel or carrying accessories.
    - Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
    - o Baby weight scales.
    - o Garments or other products that allow hands-free pump operation.
    - o Breast milk storage bags, ice-packs, labels, labeling lids, and other similar products.
    - o Nursing bras, bra pads, breast shells, nipple shields, and other similar products.
    - Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

**Note**: See the <u>Indications for Coverage</u> section above for covered breastfeeding equipment.

## TRAVEL IMMUNIZATIONS: ADDITIONAL INFORMATION

Immunizations that are specific to **travel** (e.g., typhoid, yellow fever, cholera, plague, and Japanese encephalitis virus) are not required by PPACA and are excluded from coverage. However, travel immunizations are available as a buy-up coverage option on certain plans. Please see the member specific plan document for details.

#### DEFINITIONS

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

**Independent Genetics Provider (for Medical Necessity Benefit Plans)**: Genetic counseling is required by an independent (not employed by a genetic testing lab) genetics provider prior to genetic testing for BRCA mutations in order to inform persons being tested about the benefits and limitations of a specific genetic test as applied to a unique

person. Genetics Providers employed by or contracted with a laboratory that is part of an Integrated health system which routinely delivers health care services beyond just the laboratory testing itself are considered independent. Genetic testing for BRCA mutations requires documentation of medical necessity by one of the following who has evaluated the member and intends to engage in post-test follow-up counseling:

- Board-eligible or Board-Certified Genetic Counselor (CGC)
- Advanced Genetics Nurse (AGN-BC)
- Genetic Clinical Nurse (GCN)
- Advanced Practice Nurse in Genetics (APNG)
- A Board-eligible or Board-Certified Clinical Geneticist
- A Board-certified physician with experience in cancer genetics (defined as providing cancer risk assessment on a regular basis and having received specialized ongoing training in cancer genetics.)

**Modifier 33**: Preventive service; when the primary purpose of the service is the delivery of an evidence based service in accordance with a US Preventive Services Task Force A or B rating in effect and other preventive services identified in preventive services mandates (legislative or regulatory), the service may be identified by adding 33 to the procedure. For separately reported services specifically identified as preventive, the modifier should not be used.

Please see <u>Applicable Codes</u> section below for more information about Modifier 33.

#### APPLICABLE CODES

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this guideline does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Coverage Determination Guidelines may apply.

## Modifier 33

UnitedHealthcare considers the procedures and diagnostic codes and Preventive Benefit Instructions listed in the table below in determining whether preventive care benefits apply. While modifier 33 may be reported, it is not used in making preventive care benefit determinations.

CPT®	is a	registered	trademark of	the American	Medical Association	on

## **Preventive Care Services**

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Abdominal Aortic Aneurysm Screening <u>USPSTF Rating (June 2014): B</u> The USPSTF recommends one- time screening for abdominal aortic aneurysm (AAA) by ultrasonography in men ages 65 to 75 years who have ever smoked.	<ul> <li>Procedure Code(s): Ultrasound screening study for abdominal aortic aneurysm: • 76706</li> <li>Diagnosis Code(s): • <u>ICD-10</u>: Z87.891, F17.210, F17.211, F17.213, F17.218, F17.219</li> </ul>	<ul> <li>Age 65 through 75 (ends on 76<sup>th</sup> birthday)</li> <li>One of the Diagnosis Codes listed in this row.</li> </ul>
Bacteriuria Screening <u>USPSTF Rating (July 2008): A</u> Screening for asymptomatic bacteriuria with urine culture for pregnant women at 12 to 16 weeks' gestation or at the first prenatal visit, if later.	<ul> <li>Procedure Code(s):</li> <li>81007</li> <li>Diagnosis Code(s):</li> <li>Pregnancy Diagnosis Code (see list at end of section).</li> </ul>	<ul> <li>Payable with a Pregnancy Diagnosis Code (see list at end of section)</li> </ul>

Preventive Care Services

UnitedHealthcare Commercial Coverage Determination Guideline

Proprietary Information of UnitedHealthcare. Copyright 2017 United HealthCare Services, Inc.

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventiv	e care medications refer to pharr	nacy plan automistrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
<ul> <li>Chlamydia Infection Screening</li> <li>USPSTF Rating (Sept. 2014): B The USPSTF recommends screening for chlamydia in sexually active women age 24 years and younger and in older women who are at increased risk for infection.</li> <li>Note: This recommendation applies to all sexually active adolescents and adult women, including pregnant women.</li> <li>Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11 years – 21 years.</li> <li>Gonorrhea Screening</li> <li>USPSTF Rating (Sept. 2014): B The USPSTF recommends screening for gonorrhea in sexually active women age 24 years and younger and in older women who are at increased risk for infection.</li> <li>Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11 years – 21 years.</li> </ul>	<ul> <li>Procedure Code(s): Chlamydia Infection Screening:</li> <li>86631, 86632, 87110, 87270, 87320, 87490, 87491, 87492, 87801, 87810</li> <li>Blood draw:</li> <li>36415, 36416 Blood draw codes only apply to lab codes 86631 or 86632.</li> <li>Diagnosis Code(s): Pregnancy:</li> <li>Pregnancy Diagnosis Code (see list at end of section), OR Screening: <u>ICD-10</u>:</li> <li>Adult: Z00.00, Z00.01</li> <li>Child: Z00.121, Z00.129</li> <li>Other:Z11.3, Z11.8, Z11.9, Z20.2</li> <li>Procedure Code(s): 87590, 87591, 87592, 87801, 87850</li> <li>Diagnosis Code(s): Pregnancy:</li> <li>Pregnancy Diagnosis Code (see list at end of section), OR Screening:</li> <li><u>CD-10</u>:</li> <li>Adult:Z00.00, Z00.01</li> <li>Child: Z00.121, Z00.129</li> <li>Other:Z11.3, Z11.9, Z20.2</li> </ul>	<ul> <li>Chlamydia Infection Screening:</li> <li>Payable with a Pregnancy Diagnosis Code (see list at end of section) OR</li> <li>One of the Screening Diagnosis Codes listed in this row.</li> <li>Blood draw: <ul> <li>Payable when billed with both of the following:</li> <li>With 86631 or 86632 <u>AND</u></li> </ul> </li> <li>With one of the Screening Diagnosis Codes listed in this row OR with a Pregnancy Diagnosis Code (see list at end of section)</li> <li>Payable with either a Pregnancy Diagnosis Code (see list at end of section) OR</li> <li>One of the Screening Diagnosis Codes listed in this row.</li> </ul>
Hepatitis B Virus Infection ScreeningPregnant Women: USPSTF Rating (June 2009): A Screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit.Persons at High Risk: USPSTF Rating (May 2014): B The USPSTF recommends screening for hepatitis B virus	<ul> <li>Procedure Code(s): Hepatitis B Virus Infection Screening: <ul> <li>87340, 87341, G0499</li> </ul> </li> <li>Blood draw: <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): Pregnancy: <ul> <li>Pregnancy:</li> <li>Pregnancy Diagnosis Code (see list at end of section), OR</li> </ul> </li> <li>Screening: <ul> <li>ICD-10: Z57.8, Z00.00, Z00.01, Z11.59</li> </ul> </li> </ul>	<ul> <li>Hepatitis B Virus Infection Screening <ul> <li>Payable with a Pregnancy Diagnosis Code (see list at end of section) <b>OR</b></li> <li>One of the Screening Diagnosis Codes listed in this row.</li> </ul> </li> <li>Blood draw: <ul> <li>Payable when billed with one of the listed Hepatitis B Virus Infection Screening procedure codes listed in this row <b>AND</b></li> <li>With a Pregnancy Diagnosis Code (see list at end of section) <b>OR</b> one of the Screening Diagnosis Codes listed in this row.</li> </ul> </li> </ul>

Preventive Care Services Page 7 of 50 UnitedHealthcare Commercial Coverage Determination Guideline Effective 10/01/2017 Proprietary Information of UnitedHealthcare. Copyright 2017 United HealthCare Services, Inc.

Tor prevention	e care medications refer to phari	nacy plan aunimistrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
(HBV) infection in <b>persons at</b> <b>high risk</b> for infection.		
Please also see Medical Policy: <u>Hepatitis Screening</u>		
Hepatitis C Virus Infection Screening <u>USPSTF Rating (June 2013): B</u> The USPSTF recommends screening for hepatitis C virus (HCV) infection in persons at high risk for infection. The USPSTF also recommends offering one-time screening for HCV infection to adults born between 1945 and 1965. Please also see Medical Policy:	<ul> <li>Procedure Code(s): Hepatitis C Virus Infection Screening:</li> <li>86803, 86804, G0472</li> <li>Blood draw:</li> <li>36415, 36416</li> <li>Diagnosis Code(s):</li> <li>Hepatitis C Virus Infection Screening Diagnosis Code (see list at end of section.)</li> </ul>	<ul> <li>Hepatitis C Virus Infection Screening:</li> <li>Preventive with one of the Hepatitis C Virus Infection Diagnosis codes (see list at the end of section).</li> <li>Blood draw:</li> <li>Preventive with one of the Hepatitis C Virus Infection Screening procedure codes listed in this row AND a Hepatitis C Virus Infection Screening Diagnosis Code (see list at end of section).</li> </ul>
Hepatitis Screening		
<ul> <li>HIV - Human Immunodeficiency Virus - Screening for Adolescents and Adults</li> <li>USPSTF Rating (April 2013): A <ul> <li>The USPSTF recommends that clinicians screen for HIV infection in adolescents and adults ages 15 to 65 years. Younger adolescents and older adults who are at increased risk should also be screened.</li> <li>The USPSTF recommends that clinicians screen all pregnant women for HIV, including those who present in labor who are untested and whose HIV status is unknown.</li> </ul> </li> <li>Note: Bright Futures recommends HIV screening lab work be conducted once between ages 15 years - 18 years. Also recommended anytime between ages 11 - 14 years, and 19 - 21 years when a</li> </ul>	Procedure Code(s): <i>HIV – Human Immunodeficiency</i> <i>Virus – Screening:</i> • 86689, 86701, 86702, 86703, 87389, 87390, 87391, 87806, G0432, G0433, G0435, G0475, S3645 <i>Blood draw:</i> • 36415, 36416 <b>Diagnosis Code(s):</b> <i>Pregnancy:</i> • Pregnancy Diagnosis Code (see list at end of section) <b>OR</b> <i>Screening:ICD-10:</i> • Adult: Z00.00, Z00.01 • Child: Z00.121, Z00.129, • Other: Z22.6, Z22.8, Z22.9, Z11.3, Z11.4, Z11.59, Z11.9, Z20.6 <i>Also see Expanded Women's</i> <i>Preventive Health table below.</i>	<ul> <li>No age limits.</li> <li>HIV - Human Immunodeficiency Virus - Screening:</li> <li>Preventive when billed with a Pregnancy Diagnosis Code (see list at end of section) OR</li> <li>One of the Screening Diagnosis Codes listed in this row.</li> <li>Blood draw:</li> <li>Payable when billed with both of the following: <ol> <li>With one of the listed HIV Screening procedure codes listed in this row</li> </ol> </li> <li>AND</li> <li>With one of the following: <ol> <li>one of the Screening Diagnosis Codes listed in this row, OR</li> <li>with a Pregnancy Diagnosis Code (see list at end of section)</li> </ol> </li> </ul>

## Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

i el preventiv	e care medications refer to pharm	nacy plan administrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
RH Incompatibility Screening <u>USPSTF Rating (Feb. 2004): A</u> Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care. <u>USPSTF Rating (Feb. 2004): B</u> Repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.	<ul> <li>Procedure Code(s): <i>RH Incompatibility Screening:</i> <ul> <li>86901</li> </ul> </li> <li><i>Blood draw:</i> <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>Pregnancy Diagnosis Code (see list at end of section).</li> </ul> </li> </ul>	<ul> <li><i>RH Incompatibility Screening:</i></li> <li>Payable with a Pregnancy Diagnosis Code (see list at end of section)</li> <li><i>Blood draw:</i></li> <li>Payable when billed with 86901 <u>AND</u> with a Pregnancy Diagnosis Code (see list at end of section)</li> </ul>
Syphilis Screening <u>USPSTF Rating (June 2016): A</u> The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection. (Asymptomatic, nonpregnant adults and adolescents who are at increased risk for syphilis infection) <u>USPSTF Rating (May 2009): A</u> The USPSTF recommends that clinicians screen all pregnant women for syphilis infection.	<ul> <li>Procedure Code(s): Syphilis Screening: • 86592, 86593</li> <li>Blood draw: • 36415, 36416</li> <li>Diagnosis Code(s): Pregnancy: • Pregnancy Diagnosis Code (see list at end of section) <u>OR</u> Screening: • <u>ICD-10:</u> Z00.00, Z00.01, Z00.121, Z00.129, Z11.2, Z11.3, Z11.9, Z20.2</li> </ul>	<ul> <li>Syphilis Screening:</li> <li>Payable with a Pregnancy Diagnosis Code (see list at end of section) OR</li> <li>One of the Screening Diagnosis Code listed in this row.</li> <li>Blood draw:</li> <li>Payable when billed with both of the following: <ol> <li>With one of the listed Syphilis Screening procedure codes listed in this row AND</li> <li>With one of the following: <ol> <li>one of the listed Screening diagnosis codes in this row OR</li> <li>with a Pregnancy Diagnosis Code(see list at end of section)</li> </ol> </li> </ol></li></ul>
Genetic Counseling and Evaluation for BRCA Testing; and BRCA Lab Screening USPSTF Rating (Dec. 2013): B The USPSTF recommends that primary care providers screen women who have family members with breast, ovarian, tubal, or peritoneal cancer with one of several screening tools designed to identify a family history that may be associated with an increased risk for potentially harmful mutations in breast cancer susceptibility genes ( <i>BRCA1</i> or <i>BRCA2</i> ). Women with positive screening results should receive genetic counseling and, if indicated after	Genetic Counseling and Evaluation: Procedure Code(s): Medical genetics and genetic counseling services: • 96040, S0265 Evaluation and Management (Office Visits): • 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, G0463 Diagnosis Code(s): • <u>ICD-10:</u> Z80.3, Z80.41, Z85.3, Z85.43, Z15.01, Z15.02	<ul> <li>Genetic Counseling and Evaluation:</li> <li>*Medical Necessity plans require genetic counseling before BRCA Lab Screening.</li> <li>Payable as preventive with one of the Genetic Counseling and Evaluation Diagnosis Codes listed in this row in primary position.</li> <li>BRCA Lab Screening:</li> <li>*Prior authorization requirements apply to BRCA lab screening.</li> <li>Payable for age 18+ when billed with one of the BRCA Lab Screening Diagnosis codes listed in this row.</li> <li>Blood draw:</li> <li>Payable when billed with both of the following:</li> </ul>

Preventive Care Services

Page 9 of 50 Effective 10/01/2017

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

i er preventit	re care medications refer to pharm	
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
counseling, BRCA testing. Please see Medical Policy titled: <u>Genetic Testing for Hereditary</u> <u>Breast Ovarian Cancer</u> <u>Syndrome (HBOC)</u>	<ul> <li>BRCA Lab Screening: Procedure Code(s):</li> <li>81211, 81212, 81213, 81214, 81215, 81216, 81217, 81162</li> <li>Blood draw:</li> <li>36415, 36416</li> <li>Diagnosis Code(s): Family History or Personal History of breast cancer and/or ovarian cancer:</li> <li><u>ICD-10</u>: Z80.3, Z80.41, Z85.3, Z85.43, Z15.01, Z15.02</li> </ul>	<ol> <li>With one of the listed BRCA Lab Screening procedure codes listed in this row, <u>AND</u></li> <li>With one of the BRCA Lab Screening diagnosis codes listed in this row.</li> </ol>
Diabetes Screening USPSTF Rating (Oct. 2015): B The USPSTF recommends screening for abnormal blood glucose as part of cardiovascular risk assessment in adults aged 40 to 70 years who are overweight or obese. Clinicians should offer or refer patients with abnormal blood glucose to intensive behavioral counseling interventions to promote a healthful diet and physical activity. See Expanded Women's Preventive Health table below for Gestational Diabetes Screening.	Procedure Code(s): Diabetes Screening: • 82947, 82948, 82950, 82951, 82952, 83036 Blood draw: • 36415, 36416 Diagnosis Code(s): REQUIRED DIAGNOSIS CODES (requires at least one): • <u>ICD-10</u> : Z00.00, Z00.01, Z13.1 AND one of the following Additional Diagnosis Codes as follows: ADDITIONAL DIAGNOSIS CODES (requires at least one): <u>OVERWEIGHT:</u> • <u>ICD-10</u> : E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29 <u>OBESITY :</u> • <u>ICD-10</u> : E66.01, E66.09, E66.1, E66.8, E66.9, Z68.41, Z68.42, Z68.43, Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39 <u>BODY MASS INDEX 40.0 AND</u> <u>OVER:</u> • <u>ICD-10</u> : Z68.41, Z68.42, Z68.43, Z68.44, Z68.45	<ul> <li>Limited to age 40-70years (ends on 71<sup>st</sup> birthday).</li> <li>Diabetes Screening: <ul> <li>Payable with one of the Required Diagnosis Codes listed in this row.</li> </ul> </li> <li>With one of the listed Additional Diagnosis Codes in this row.</li> </ul> <li>Blood draw: <ul> <li>Payable when billed with ALL of the following: <ul> <li>With one of the listed Diabetes Screening procedure codes listed in this row AND</li> <li>With one of the listed Required Diagnosis Codes AND</li> <li>With one of the listed Additional Diagnosis Codes.</li> </ul> </li> <li>Preventive Benefit Does Not Apply: <ul> <li>If a Diabetes Diagnosis Code is present in any position, the preventive benefit does not apply. See Diabetes Diagnosis Codes table below.</li> </ul> </li> </ul></li>

Preventive Care Services

Page 10 of 50 Effective 10/01/2017

Preventive Care Services				
Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines.				
	e care medications refer to pharm			
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:		
when the benefit is chective.	ESSENTIAL HYPERTENSION:			
	<ul> <li>HYPERTENSIVE HEART DISEASE:</li> <li>ICD-10: I11.0, I11.9</li> <li>HYPERTENSIVE CHRONIC KIDNEY DISEASE:</li> <li>ICD-10: I12.0, I12.9</li> <li>HYPERTENSIVE HEART AND CHRONIC KIDNEY DISEASE:</li> <li>ICD-10: I13.0, I13.10, I13.11, I13.2</li> <li>SECONDARY HYPERTENSION:</li> <li>ICD-10: I15.0, I15.1, I15.2, I15.8, I15.9, N26.2</li> <li>HYPERTENSION COMPLICATING PREGNANCY, CHILDBIRTH AND THE PUERPERIUM:</li> <li>ICD-10: 010.011, 010.012, 010.013, 010.019, 010.02, 010.113, 010.119, 010.12, 010.113, 010.211, 010.212, 010.213, 010.211, 010.212, 010.23, 010.311, 010.312, 010.313, 010.319, 010.32, 010.313, 010.319, 010.32, 010.212, 010.313, 010.319, 010.32, 010.213, 010.313, 010.319, 010.32, 010.213, 010.311, 010.312, 010.213, 0</li></ul>			
	010.33, 010.411, 010.412, 010.413, 010.419, 010.42, 010.43, 010.911, 010.912, 010.913, 010.919, 010.92, 010.93, 011.1, 011.2, 011.3,011.4, 011.5, 011.9, 013.1, 013.2, 013.3, 013.4, 013.5, 013.9, 016.1, 016.2, 016.3, 016.4, 016.5, 016.9 URGENT/EMERGENCY/CRISIS HYPERTENSION • <u>ICD-10: I16.0, I16.1, I16.9</u> See Expanded Women's Preventive Health table below			
	for Gestational Diabetes Screening.			
Gestational Diabetes Mellitus Screening	See Expanded Women's Preventive Health table below for Gestational Diabetes Screening codes.	See Expanded Women's Preventive Health table below for Gestational Diabetes Screening preventive benefit instructions.		
<u>USPSTF Rating (January 2014):</u> <u>B</u>	Diabetes Screening Codes.			

## Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

i or preventiv	e care medications refer to phari	nacy plan automistrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant women after 24 weeks of gestation.		<i>Note: This benefit applies regardless of the gestational week.</i>
<i>Also see the Diabetes Screening row above for additional diabetes screening benefits.</i>		
Rubella Screening By History of Vaccination or by Serology USPSTF Rating (1996): B Screening for rubella susceptibility by history of vaccination or by serology is recommended for all women of childbearing age at their first clinical encounter.	<ul> <li>Procedure Code(s):</li> <li>Rubella Screening by History of Vaccination: <ul> <li>No codes (included in exam)</li> </ul> </li> <li>Rubella Screening by Serology: <ul> <li>86762</li> </ul> </li> <li>Blood draw: <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>ICD-10: Z00.00, Z00.01, Z11.59, Z20.4</li> </ul> </li> </ul>	<ul> <li>Rubella Screening by Serology:</li> <li>Payable with one of the listed diagnosis codes in this row.</li> <li>Blood draw:</li> <li>Payable when billed with 86762 <u>AND</u> one of the listed diagnosis codes in this row</li> </ul>
Screening Mammography USPSTF Rating: B (2002 Recommendation) The USPSTF recommends screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older. Also, see Medical Policy titled: <u>Breast Imaging for Screening</u> and Diagnosing Cancer.	<ul> <li>Procedure Code(s): <ul> <li>G0202, 77063, 77067</li> <li>Revenue code: 0403</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul> </li> </ul>	<ul> <li>Payable regardless of age.</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li><u>Note the following:</u></li> <li>This benefit only applies to screening mammography.</li> </ul>
Cervical Cancer Screening, Pap Smear USPSTF Rating (March 2012): A Screening for cervical cancer in women ages 21 to 65 years with cytology (Pap smear) every 3 years or, for women ages 30 to 65 years who want to lengthen the screening interval, screening with a combination of cytology and human papillomavirus (HPV) testing every 5 years. Bright Futures, March 2014:	<ul> <li>Procedure Code(s):</li> <li>Code Group 1: <ul> <li>G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, Q0091, P3000, P3001</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul> </li> <li>Code Group 2: <ul> <li>88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88154, 88155, 88164, 88165, 88166, 88167, 88174, 88174, 88175</li> </ul> </li> </ul>	<ul> <li>Limited to age 21 years - 65 years (ends on 66<sup>th</sup> birthday)</li> <li>Code Group 1: <ul> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> </ul> </li> <li>Code Group 2: <ul> <li>Payable with one of the Diagnosis Codes listed in this row.</li> </ul> </li> </ul>

Preventive Care Services

Page 12 of 50 Effective 10/01/2017

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

Service: A date in this column is when the rating was released, not when the benefit is effective.         Code(s):         Preventive Benefit Instructions:           A date in this column is when the rating was released, not when the benefit is effective.         Code Group 2 Diagnosis Code(s):         Preventive Benefit Instructions:           Adolescents should no longer be routinely screening (Lipid Disorders Screening)         Code Group 2 Diagnosis Code(s):         Preventive Benefit Instructions:           Cholesterol Screening (Lipid Disorders Screening)         Procedure Code(s): Cholesterol Screening: USPSTF Rating (Nov. 2016): B Statin Use for the Primary Preventiou a fastory of Cardiovascular of Cardiovascular issease on ischemic stroke, use and mortality when all of the following circlena are met: 1) they are aged 40 to 75 years; 2) they have 1 or more CVD is greater. Identification of dysipidemia and calculation of disorders. USPSTF Rating: A Screening men aged 35 and older for lipid disorders. USPSTF Rating: A Screening men aged 35 and older for lipid disorders. USPSTF Rating: A Screening men aged 35 and older for lipid disorders. USPSTF Rating: A Screening women aged 45 and ulters for finid disorders if they are at increased risk for coromary heart disease. USPSTF Rating: A Screening women aged 45 and ulters for finid disorders if they are at increased risk for coromary heart diseases.         Screen	A date in this column is when the rating was released, not when the benefit is effective.       Code Group 2 Diagnosis Code(5): . If LD-10: 000 00, 200 01, 201411, 201419, 212.4         Adolescents should no longer be routinely screened for cervical dysplasia until age 21.       Code Group 2 Diagnosis Code(5): . If LD-10: 200 00, 200 01, 201411, 201419, 212.4         Cholesterol Screening)       Proceeding Code(5): Cholesterol Screening: . 83061, 82456, 83718, 83719, 83721, 84478       Age 40 – 75 years (ands on 76 <sup>th</sup> birthday)         USPSTF Rating (Nov. 2016): B Issaein Adults - The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (ie, symptomatic coronary after following criteria are met: 1) they have 1 or more CVD risk factors (ie, dyslipidemi, diabetes, hypertension, or smaking): and at calculation of dyslipidemia and calculation of dyslipidemia and calculation of dyslipidemia and calculation of lo-year CVD events and mortality screening in adults aged 40 to 75 years.       Additional Diagnosis Codes: see Preventive Benefit Instrumer Strate (IE) (IE) (IE) (IE) (IE) (IE) (IE) (IE)	For preventiv	e care medications refer to phari	macy plan administrator.
routinely screened for cervical dysplasia until age 21.       Code Group 2 Diagnosis (CD2-12; 200.00, 200.01, 201.411, 201.419, 201.419, 201.419, 201.419, 201.419, 201.419, 201.419, 201.419, 201.419, 201.419, 201.411, 201.411, 201.41	routinely screened for cervical dysplasia until age 21.Code Group 2 Diagnosis (201411, 201419, 212.4)Cholesterol Screening (Lipid Disorders Screening)Procedure Code(s): (201411, 201419, 212.4)Age 40 – 75 years (ends on 76th birthday)USPSTF Rating (Nov, 2016): B Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (ie, teast one): • 102-101; 200.00, 200.01, Z13.220Age 40 – 75 years (ends on 76th birthday) • Preventive with one of the Required Diagnosis Codes: (articestore): 2016211; 200.00, 200.01, Z13.220Iber brinders (articestore): (articestore): (articestore): 201671; 201.00, 200.01, Z13.220Age 40 – 75 years (ends on 76th bir now.Biood draw: • 36415, 36416Biaonasis Codes: (articestore): 200.00, 200.01, Z13.220Biood draw: • 102-101; 200.00, 200.01, Z13.220Iber and mortality when all of the Primary (articestore): 2016; 272.0, 282.49, (articestore): 272.0, 282.49, 283.42, 287.898.171.2218, • 102-101; 272.0, 282.49, 283.42, 287.898.171.2218, • 102-102; 125.66.01, E66.09, E66.1, E66.8, E66.9Biod draw: • Payable for age 25 - 39 years and age 76 years and dider when billed with one of the Required Diagnosis Codes listed in this row.Biood draw: • 102-102; 272.0, 282.49, 283.42, 287.898.41, 268.42, 266.43, 268.41, 268.45, • 102-102; 110Biood draw: • Payable for age 25 - 39 years and age 76 years and dider when billed with one of the Required Diagnosis Codes listed in this row.Biood draw: • 102-102; 125.66.01, E66.09, 10-year CVD event risk requireds • 102-102; 110, 56.11, E66.9, 10-year CVD event risk	A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Disorders Screening)Cholesterol Screening: • Preventive with one of the Required Diagnosis Codes listed in this row.USPSTF Rating (Nov. 2016): B Statin Use for the Primary Disease in Adults - The USPST Frecommends that adults without a history of cardiovascular disease (CVD) (its, symptomatic coronary artery disease or ischemic strokenic stroken	Disorders Screening)Cholesterol Screening: Biodod raw:• Preventive with one of the Required Diagnosis Codes listed in this row.USESTF Rating (Nov. 2016): B Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (ie, symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the preventive of CVD events and mortality when all of the Following circle are met: 1) they have 1 or more CVD risk factors (ie, dyslipidemia, diabets, hypertension, or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipid screening in administrator.Cholesterol Screening Men (June 2008)*: . (CD-10: 272.0, 282.49, 283.42, 287.891, F17.210, F17.211, F17.213, F17.218, F17.211, F17.213, F17.218, F17.211, F17.213, F17.218, F17.211, F17.213, F17.219, F17.211, F17.213, F17.218, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.210, F17.211, F17.213, F17.210, F17.211, F17.213, F17.210, F17.211, F17.213, F17.218, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.219, F17.211, F17.213, F17.210, F17.211, F17.213, F1	routinely screened for cervical	Code(s): • <u>ICD-10:</u> Z00.00, Z00.01,	
		Disorders Screening) USPSTF Rating (Nov. 2016): B Statin Use for the Primary Prevention of Cardiovascular Disease in Adults – The USPSTF recommends that adults without a history of cardiovascular disease (CVD) (ie, symptomatic coronary artery disease or ischemic stroke) use a low- to moderate-dose statin for the prevention of CVD events and mortality when all of the following criteria are met: 1) they are aged 40 to 75 years; 2) they have 1 or more CVD risk factors (ie, dyslipidemia, diabetes, hypertension, or smoking); and 3) they have a calculated 10-year risk of a cardiovascular event of 10% or greater. Identification of dyslipidemia and calculation of 10-year CVD event risk requires universal lipids screening in adults aged 40 to 75 years. Note: for statin medications benefits refer to pharmacy plan administrator. Screening Men (June 2008)*: • Screening men aged 35 and older for lipid disorders. USPSTF Rating: A • Screening men aged 20 to 35 for lipid disorders if they are at increased risk for coronary heart disease. USPSTF Rating: B Screening Women at Increased Risk (June 2008)*: • Screening women aged 45 and older for lipid disorders of reconary heart disease. Screening women aged 45 and older for lipid disorders if	Procedure Code(s): Cholesterol Screening: • 80061, 82465, 83718, 83719, 83721, 84478 Blood draw: • 36415, 36416 Diagnosis Code(s): Required Diagnosis Codes: (at least one): • <u>ICD-10</u> ; Z00.00, Z00.01, Z13.220 Additional Diagnosis Codes: See Preventive Benefit Instructions Column For When Required: FAMILY or PERSONAL HISTORY: • <u>ICD-10</u> ; Z72.0, Z82.49, Z83.42, Z87.891, F17.210, F17.211, F17.213, F17.218, F17.219 OBESITY: • <u>ICD-10</u> ; E66.01, E66.09, E66.1, E66.8, E66.9 BODY MASS INDEX 40 AND OVER, ADULT: • <u>ICD-10</u> ; Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 ESSENTIAL HYPERTENSION: • <u>ICD-10</u> ; I10 SECONDARY HYPERTENSION: • <u>ICD-10</u> ; I15.0, I15.1, I15.2, I15.8, I15.9, N26.2 HYPERTENSION COMPLICATING PREGNANCY, CHILDBIRTH AND THE PUERPERIUM: • <u>ICD-10</u> ; O10.011, O10.012, O10.013, O10.019, O10.02,	<ul> <li>Preventive with one of the Required Diagnosis Codes listed in this row.</li> <li>Blood draw:         <ul> <li>Payable for age 40 – 75 years when billed with one of the listed Cholesterol Screening procedure codes AND with one of the Required Diagnosis Codes listed in this row.</li> </ul> </li> <li>Age 35 – 39 years, and age 76 years and older:         <ul> <li>Preventive with one of the Required Diagnosis Codes listed in this row.</li> </ul> </li> <li>Blood draw:         <ul> <li>Payable for age 35 – 39 years and age 76 years and older:</li> <li>Preventive with one of the Required Diagnosis Codes listed in this row.</li> </ul> </li> <li>Blood draw:         <ul> <li>Payable for age 35 – 39 years and age 76 years and older when billed with one of the listed Cholesterol Screening procedure codes AND with one of the Required Diagnosis Codes listed in this row.</li> </ul> </li> <li>Age 20 to 34 (ends on 35<sup>th</sup> birthday):         <ul> <li>Preventive with one of the Required Diagnosis Codes listed in this row AND with one of the listed Additional Diagnosis Codes listed in this row AND with one of the listed Cholesterol Screening procedure codes AND with one of the listed Cholesterol Screening procedure codes AND             <ul> <li>with one of the listed Cholesterol Screening procedure codes AND</li> <li>with one of the Required Diagnosis Codes listed in this row AND</li> <li>with one of the Required Diagnosis Codes listed in this row AND</li> <li>with one of the Required Diagnosis Codes listed in this row AND</li> <li>with one of the Required Diagnosis Codes listed in this row AND</li> <li>with one of the Required Diagnosis Codes listed in this row AND</li> <li>with one of the Required Diagnosis Codes listed in this row</li> </ul> </li></ul></li></ul>

Preventive Care Services

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

Tor preventiv	e care medications refer to phari	nacy plan automistrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
coronary heart disease. USPSTF Rating: A • Screening women aged 20 to 45 for lipid disorders if they are at increased risk for coronary heart disease. USPSTF Rating: B *Preventive coverage for the 2008 ratings expire Nov. 30, 2017 *Note: See Dyslipidemia Screening (Bright Futures) row for recommendations for children.	<ul> <li>010.13, 010.211, 010.212, 010.213, 010.219, 010.22, 010.23, 010.311, 010.312, 010.313, 010.319, 010.32, 010.33, 010.411, 010.412, 010.413, 010.419, 010.42, 010.43, 010.911, 010.912, 010.93, 010.919, 010.92, 010.93, 011.1, 011.2, 011.3, 011.4, 011.5, 011.9, 013.1, 013.2, 013.3, 013.4, 013.5, 013.9, 016.1, 016.2, 016.3, 016.4, 016.5, 016.9</li> <li>URGENT/EMERGENCY/CRISIS HYPERTENSION</li> <li>ICD-10: I16.0, I16.1, I16.9</li> <li>DIABETES:</li> <li>See Diabetes Diagnosis Code List table below.</li> <li>ATHERO-SCLEROSIS:</li> <li>See Atherosclerosis Diagnosis Code List table below</li> <li>CORONARY ATHERO-SCLEROSIS:</li> <li>ICD-10: I25.10, I25.110, I25.711, I25.118, I25.119, I25.709, I25.710, I25.711, I25.710, I25.710, I25.711, I25.718, I25.719, I25.720, I25.721, I25.728, I25.729, I25.730, I25.731, I25.738, I25.739, I25.750, I25.751, I25.758, I25.759, I25.760, I25.761, I25.768, I25.769, I25.790, I25.711, I25.798, I25.790, I25.711, I25.798, I25.799, I25.810, I25.811, I25.812</li> </ul>	apply: • <u>ICD-10:</u> E71.30, E75.5, E78.00, E78.01, E78.2, E78.3, E78.4, E78.5, E78.79, E78.81, E78.89, E88.2, E88.89
Colorectal Cancer Screening USPSTF Rating (June 2016): A The USPSTF recommends	Fecal Occult Blood Testing (FOBT), Fecal Immunochemical Test (FIT), Fecal DNA, Sigmoidoscopy, or	Age Limits for Colorectal Cancer Screenings: • 50 years – 75 years (ends on 76 <sup>th</sup> birthday)
screening for colorectal cancer starting at age 50 years and continuing until age 75 years. The risks and benefits of different screening methods vary.	Procedure Code(s): Colonoscopy: Procedure Code(s): Code Group 1: • Sigmoidoscopy: G0104, G0106 • Colonoscopy: G0105, G0120, G0121, G0122 • FOBT and FIT: G0328 • Colonoscopy Pre-op	Fecal Occult Blood Testing (FOBT), Fecal Immunochemical Test (FIT), Fecal DNA, Sigmoidoscopy, or Colonoscopy: <u>Code Group 1</u> : Does not have diagnosis code requirements for preventive benefits to apply.

Preventive Care Services

# Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

	e care medications refer to phari	
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
	Consultation: S0285	<u>Code Group 2</u> : Paid as preventive if:
	Consultation: S0285 Code Group 2: • Sigmoidoscopy: 45330, 45331, 45333, 45338, 45346 • Colonoscopy: 44388, 44389, 44392, 44394, 45378, 45380, 45381, 45384, 45385, 45388 • FOBT and FIT: 82270, 82274 Code Group 3: • Pathology: 88304, 88305 Code Group 4: • Anesthesia: 00810, 99152, 99153, 99156, 99157, G0500 Code Group 5: Pre-op/Consultation: • 99201, 99202, 99203, 99204, 99205 • 99211, 99212, 99213, 99214, 99245 • 99241, 99242, 99243, 99244, 99245 Code Group 6: Fecal DNA: 81528 Diagnosis Code(s) (for Code Group 2, 3, and 4): • ICD-10: Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z80.0, Z83.71, Z83.79 (for Code Group 5):	<ul> <li>Code Group 2: Paid as preventive if:</li> <li>billed with one of the Diagnosis Codes listed in this row OR</li> <li>billed with one of the Procedure Codes from Code Group 1, regardless of diagnosis.</li> <li>Code Group 3 (pathology) AND Code Group 4 (anesthesia):</li> <li>Paid as preventive if:</li> <li>billed with one of the Diagnosis Codes listed in this row AND</li> <li>billed with one of the Procedure Codes from Code Group 1 or Code Group 2.</li> <li>Code Group 3 and 4: Note the following: <ul> <li>Preventive when performed for a colorectal cancer screening.</li> <li>Preventive benefits only apply when the surgeon's claim is preventive.</li> </ul> </li> <li>Code Group 5 : Paid as Preventive if billed with one of the Code Group 5 diagnosis codes.</li> <li>Code Group 6 (Fecal DNA):</li> <li>Benefit is limited to once every 3 years.</li> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> </ul>
	• <u>ICD-10:</u> Z12.10, Z12.11, Z12.12, Z80.0, Z83.71, Z83.79 Computed Tomographic Colonography (Virtual	<ul> <li>Computed Tomographic Colonography (Virtual Colonoscopy):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li>Prior authorization requirements may</li> </ul>
	<ul> <li>Colonoscopy):</li> <li>Procedure Code(s):</li> <li>74263</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>	<ul> <li>Prior authorization requirements may apply, depending on plan.</li> </ul>
Wellness Examinations (well baby, well child, well adult) <u>USPSTF Rating: None</u> UHC supports AAP and AAFP age and frequency guidelines.	<ul> <li>Procedure Code(s):</li> <li>G0402, G0438, G0439</li> <li>G0445, S0610, S0612, S0613</li> <li>99381, 99382, 99383, 99384, 99385, 99386, 99387</li> <li>99391, 99392, 99393, 99394, 99395, 99396, 99397</li> </ul>	<ul> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li>G0445 is limited to twice per year.</li> <li>G0296 is limited to age 55 to 80 years (ends on 81<sup>st</sup> birthday)</li> </ul>

Preventive Care Services

#### Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
<ul> <li>HHS Requirements:</li> <li>These codes also include the following HHS requirements for Women:</li> <li>Breastfeeding support and counseling</li> <li>Contraceptive methods counseling</li> <li>Domestic violence screening</li> <li>Annual HIV counseling</li> <li>Sexually Transmitted Infections counseling</li> <li>Well-woman visits</li> </ul>	<ul> <li>99401, 99402, 99403, 99404</li> <li>99411, 99412</li> <li>99461</li> <li>Counseling Visit (to Discuss the Need for Lung Cancer Screening (LDCT) Using Low Dose CT Scan):</li> <li>G0296</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li>Also see Expanded Women's Preventive Health table</li> </ul>	
	below.	
Immunizations <u>USPSTF Rating: None</u> An immunization that does not fall under one of the exclusions in the Certificate of Coverage is considered covered after the following conditions are satisfied: (1) FDA approval; (2) explicit ACIP recommendation published in the Morbidity & Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC). Implementation will typically occur within 60 days after publication in the MMWR.	See Preventive Immunizations table below.	See Preventive Immunizations table below.
Newborn Screenings All newbornsUSPSTF Rating (July 2008): B Hearing Screening - screening for hearing loss in all newborn infantsUSPSTF Rating (March 2008): A Hypothyroidism Screening - screening for congenital hypothyroidism in newbornsUSPSTF Rating (March 2008): A Phenylketonuria Screening - screening for phenylketonuria (PKU) in newborns	Procedure Code(s): Hearing Screening: • V5008, 92551, 92558, 92585, 92586, 92587, 92588 Hypothyroidism Screening: • 84437, 84443 <i>Blood draw:</i> • 36415, 36416 Phenylketonuria Screening: • S3620, 84030 <i>Blood draw:</i> • 36415, 36416 Sickle Cell Screening: • S3850, 83020, 83021, 83030,	<ul> <li>Newborn Screenings:</li> <li>Age 0 – 90 days</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li>Blood draw:</li> <li>Age 0-90 days, payable when billed with one of the listed Hypothyroidism Screening, Phenylketonuria Screening, or Sickle Cell Screening procedure codes.</li> </ul>

Preventive Care Services

#### Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

i or preventiv	For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:	
USPSTF Rating (Sept. 2007): A Sickle Cell Screening - screening for sickle cell disease in newborns Note: For Bright Futures hearing screening, see row below Hearing Tests (Bright Futures).	<ul> <li>83033, 83051</li> <li>Blood draw:</li> <li>36415, 36416</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>		
Metabolic Screening Panel (newborns)	<ul> <li>Procedure Code(s): <i>Metabolic Screening Panel:</i> <ul> <li>\$3620, 82017, 82136, 82261, 82775, 83020, 83498, 83516, 84030, 84437, 84443</li> </ul> </li> <li><i>Blood draw:</i> <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul> </li> </ul>	<ul> <li>Metabolic Screening Panel:</li> <li>Age 0 – 90 days</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> <li>Blood draw:</li> <li>Age 0-90 days, payable when billed with one of the listed Metabolic Screening Panel Procedure Codes listed in this row.</li> </ul>	
Osteoporosis Screening <u>USPSTF Rating (Jan. 2011): B</u> The USPSTF recommends screening for osteoporosis in women age 65 and older, and in younger women whose fracture risk is equal to or greater than that of a 65-year-old white woman who has no additional risk factors.	Procedure Code(s): <ul> <li>76977, 77078, 77080, 77081</li> <li>G0130</li> </ul> Diagnosis Code(s): <ul> <li><u>ICD-10</u>: Z00.00, Z00.01, Z13.820, Z82.62</li> </ul>	<ul> <li>Preventive with one of the Diagnosis Codes listed in this row.</li> </ul>	
Prostate Cancer Screening <u>USPSTF Rating (2008): I</u> At the time PPACA was implemented the USPSTF concluded that the current evidence was insufficient to assess the balance of benefits and harms of prostate cancer screening in men younger than age 75 years.	Procedure Code(s): <u>Code Group 1:</u> • G0102, G0103 <i>Blood draw:</i> • 36415, 36416 <u>Code Group 2:</u> (requires diagnosis code) • 84152, 84153, 84154 <i>Blood draw:</i> • 36415, 36416	<ul> <li><u>Code Group 1</u>:</li> <li>Age 40 or older.</li> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> <li><u>Blood draw:</u></li> <li>Payable when billed with G0103</li> <li><u>Code Group 2</u>: Age 40 or older when billed with one of the Diagnosis Codes listed in this row.</li> </ul>	
<u>USPSTF Rating (May 2012): D</u> The USPSTF recommends against prostate specific antigen (PSA) based screening for prostate cancer. NOTE: While this screening is	<ul> <li>Diagnosis Code(s): <u>Code Group 1:</u></li> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> <li><u>Code Group 2:</u></li> <li><u>ICD-10:</u> Z00.00, Z00.01,</li> </ul>	<ul> <li>Pow.</li> <li>Blood draw:</li> <li>Payable when billed: <ul> <li>with 84152, 84153 or 84154 AND</li> <li>with one of the Diagnosis Codes listed in this row</li> </ul> </li> </ul>	

Preventive Care Services

Page 17 of 50 Effective 10/01/2017

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
not recommended by the USPSTF, the American Urological Association recommends PSA screening, together with digital rectal examination, only after explanation of the possible advantages and harms of such screening.	Z12.5, Z80.42	
Screening and Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse USPSTF Rating (May 2013): B The USPSTF recommends that clinicians screen adults aged 18 years or older for alcohol misuse and provide persons engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce alcohol misuse.	<ul> <li>Procedure Code(s): Alcohol or drug use screening:</li> <li>99408, 99409 Annual alcohol screening:</li> <li>G0442 Brief counseling for alcohol:</li> <li>G0443</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>	<ul> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> </ul>
Bright Futures (April 2017): Bright Futures recommends alcohol or drug use assessments from age 11 years – 21 years.		
<ul> <li>High Blood Pressure in Adults - Screening: USPSTF Rating (Oct. 2015):A</li> <li>The USPSTF recommends screening for high blood pressure in adults aged 18 years or older.</li> <li>The USPSTF recommends obtaining measurements outside of the clinical setting for diagnostic confirmation before starting treatment.</li> </ul>	Blood Pressure Measurement in a Clinical Setting: • n/a Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting): Procedure Code(s): Ambulatory Blood Pressure Measurement: • 93784, 93786, 93788 or 93790 Diagnosis Code: Abnormal blood-pressure reading without diagnosis of hypertension: • <u>ICD-10</u> : R03.0	<ul> <li>Blood Pressure Measurement in a Clinical Setting: <ul> <li>This service is included in a preventive care wellness examination.</li> </ul> </li> <li>Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting): <ul> <li>Age 18 years and up.</li> <li>Payable as preventive when billed with the Diagnosis Code listed in this row</li> </ul> </li> </ul>
Chemoprevention of Breast Cancer (Counseling) USPSTF Rating (July 2002): B The USPSTF recommends that clinicians discuss chemoprevention with women at high risk for breast cancer and at	Procedure Code(s): Evaluation and Management (Office Visits): • 99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397, G0463	<ul> <li>Payable as preventive when billed with one of the Diagnosis Codes listed in this row in the primary position.</li> </ul>

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
low risk for adverse effects of chemoprevention. Clinicians should inform patients of the potential benefits and harms of chemoprevention.	Diagnosis Code(s): • <u>ICD-10:</u> Z80.3, Z80.41, Z15.01, Z15.02	
Primary Care Interventions to Promote Breastfeeding USPSTF Rating (Oct. 2016): B The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.	<ul> <li>n/a</li> <li>Also see Expanded Women's</li> <li>Preventive Health table</li> <li>below.</li> </ul>	<ul> <li>Included in primary care or OB/GYN office visits.</li> </ul>
Screening for Depression in Adults <u>USPSTF Rating (Jan. 2016): B</u> Recommendation: The USPSTF recommends screening for depression in the general adult population, including pregnant and postpartum women. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up. (B recommendation)	Procedure Code(s): • 96127, G0444 Diagnosis Code(s): Required for 96127 only: • <u>ICD-10:</u> Z13.89	One of the Diagnosis Codes listed in this row is required for 96127. The Diagnosis Codes listed in this row are <b>not</b> required for G0444.
Depression in Children and Adolescents (Screening) USPSTF Rating (Feb. 2016): B The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12 to 18 years. Screening should be implemented with adequate systems in place to ensure accurate diagnosis, effective treatment, and appropriate follow-up. Note: The Bright Futures Periodicity Schedule recommends depression screening begin at age 12 through 21 years.	<pre>Procedure Code(s):     96127, G0444 Diagnosis Code(s): Required for 96127 only:     ICD-10: Z13.89</pre>	One of the Diagnosis Codes listed in this row is required for 96127. The Diagnosis Codes listed in this row are <u>not</u> required for G0444.
Behavioral Counseling in Primary Care to Promote a Healthful Diet and Physical Activity for Cardiovascular	<b>Procedure Code(s):</b> <i>Medical Nutrition Therapy or</i> <i>Counseling:</i> • 97802, 97803, 97804, G0270,	<ul> <li>G0446 is limited to once per year.</li> <li><u>One of the Diagnosis Codes listed in this row</u> are:</li> </ul>

	e care medications refer to pharm	
<b>Service:</b> A date in this column is when	Code(s):	Preventive Benefit Instructions:
the rating was released, not		
when the benefit is effective.		
Disease Prevention in Adults with Cardiovascular Risk Factors USPSTF Rating (Aug. 2014): B The USPSTF recommends offering or referring adults who are overweight or obese and	G0271, S9470 Preventive Medicine Individual Counseling: • 99401, 99402, 99403, 99404 Behavioral Counseling or Therapy: • G0446, G0447, G0473, 0403T <b>Diagnosis Code(s):</b> SCREENING:	<ul> <li>Required for 97802-97804, 99401-99404, G0270, G0271,S9470 and 0403T</li> <li><u>NOT</u> required for G0446, G0447 and G0473</li> </ul>
have additional cardiovascular disease (CVD) risk factors to intensive behavioral counseling interventions to promote a healthful diet and physical activity for CVD prevention.	<ul> <li>• <u>ICD-10</u>: Z13.220</li> <li>HISTORY:</li> <li>• <u>ICD-10</u>: Z72.0, Z87.891, Z82.49, F17.210, F17.211, F17.213, F17.218, F17.219</li> </ul>	
	OVERWEIGHT: • <u>ICD-10:</u> E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29	
	BODY MASS INDEX 30.0 – 39.9: • <u>ICD-10:</u> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39	
	BODY MASS INDEX 40.0 AND OVER: • <u>ICD-10:</u> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45	
	IMPAIRED FASTING GLUCOSE: • <u>ICD-10</u> : R73.01	
	METABOLIC SYNDROME: • <u>ICD-10</u> : E88.81	
	HYPERLIPIDEMIA / DYSLIPIDEMIA: • <u>ICD-10</u> : E78.00, E78.01, E78.1, E78.2, E78.3, E78.4, E78.5	
	OBESITY: • <u>ICD-10:</u> E66.01, E66.09, E66.1, E66.8, E66.9, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45	
	ESSENTIAL HYPERTENSION: • <u>ICD-10:</u> I10	
	SECONDARY HYPERTENSION: • <u>ICD-10:</u> I15.0, I15.1, I15.2, I15.8, I15.9, N26.2	
	HYPERTENSION COMPLICATING	

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

Service:		Preventive Benefit Instructions:
A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
	<ul> <li>PREGNANCY, CHILDBIRTH AND THE PUERPERIUM:</li> <li>ICD-10: 010.011, 010.012, 010.013, 010.019, 010.02, 010.03, 010.111, 010.112, 010.113, 010.119, 010.12, 010.13, 010.211, 010.212, 010.213, 010.219, 010.22, 010.23, 010.311, 010.312, 010.313, 010.319, 010.32, 010.33, 010.411, 010.412, 010.413, 010.419, 010.42, 010.413, 010.911, 010.912, 010.913, 010.919, 010.92, 010.93, 011.1, 011.2, 011.3, 011.4, 011.5, 011.9, 013.1, 013.2, 013.3, 013.4, 013.5, 013.9, 016.1, 016.2, 016.3, 016.4, 016.5, 016.9</li> <li>URGENT/EMERGENCY/CRISIS HYPERTENSION</li> <li>ICD-10: I16.0, I16.1, I16.9</li> </ul>	
	<ul> <li>DIABETES:</li> <li>See Diabetes Diagnosis Code List table below.</li> <li>ATHEROSCLEROSIS:</li> <li>See Atherosclerosis Diagnosis Code List table below.</li> </ul>	
	CORONARY ATHERO-SCLEROSIS: • <u>ICD-10:</u> I25.10, I25.110, I25.111, I25.118, I25.119, I25.700, I25.701, I25.708, I25.709, I25.710, I25.711, I25.718, I25.719, I25.720, I25.721, I25.728, I25.729, I25.730, I25.731, I25.738, I25.739, I25.750, I25.751, I25.758, I25.759, I25.760, I25.761, I25.768, I25.769, I25.790, I25.791, I25.798, I25.799, I25.810, I25.811, I25.812	
Screening for Obesity in Adults	Procedure Code(s): Medical Nutrition Therapy: • 97802, 97803, 97804	<ul> <li>G0446 is limited to once per year.</li> <li>One of the Diagnosis Codes listed in this row</li> </ul>
<u>USPSTF Rating (June 2012): B</u> The USPSTF recommends screening all adults for obesity. Clinicians should offer or refer	Preventive Medicine Individual Counseling: • 99401, 99402, 99403, 99404 Behavioral Counseling or Therapy:	are: • Required for 97802-97804 and 99401- 99404. • NOT required for G0446, G0447 and

Preventive Care Services

Page 21 of 50 Effective 10/01/2017

#### Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
patients with a body mass index (BMI) of 30 kg/m <sup>2</sup> or higher to intensive, multicomponent behavioral interventions.	<ul> <li>G0446, G0447, G0473</li> <li>Also see codes in the "Wellness Examinations" row above.</li> <li>Diagnosis Code(s): Body Mass Index 30.0 - 39.9: <ul> <li><u>ICD-10:</u> Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39 </li> <li>Body Mass Index 40.0 and over: <ul> <li><u>ICD-10:</u> Z68.41, Z68.42, Z68.43, Z68.44, Z68.45</li> <li>Obesity:</li> <li><u>ICD-10:</u> E66.01, E66.09, E66.1, E66.8, E66.9</li> </ul> </li> </ul></li></ul>	G0473.
Screening for Obesity in Children and Adolescents USPSTF Rating (Jan. 2010): B The USPSTF recommends that clinicians screen children aged 6 years and older for obesity and offer them or refer them to comprehensive, intensive behavioral interventions to promote improvement in weight status.	<ul> <li>Procedure Code(s): Medical Nutrition Therapy:</li> <li>97802, 97803, 97804</li> <li>Preventive Medicine Individual Counseling:</li> <li>99401, 99402, 99403, 99404</li> <li>Behavioral Counseling or Therapy:</li> <li>G0446, G0447, G0473</li> <li>Also see codes in the "Wellness Examinations" row above.</li> <li>Diagnosis Code(s): Obesity:</li> <li>ICD-10: E66.01, E66.09, E66.1, E66.8, E66.9</li> </ul>	<ul> <li>G0446 is limited to once per year.</li> <li><u>One of the Diagnosis Codes listed in this row are:</u></li> <li>Required for 97802-97804 and 99401-99404.</li> <li>NOT required for G0446, G0447 and G0473.</li> </ul>
Behavioral Counseling to Prevent Sexually Transmitted Infections <u>USPSTF Rating (Sept. 2014): B</u> The USPSTF recommends intensive behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).	<ul> <li>Procedure Code(s):</li> <li>99401, 99402, 99403, 99404</li> <li>G0445</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>	<ul> <li>G0445 is limited to twice per year.</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>
Tobacco Smoking Cessation in Adults, including Pregnant Women: Behavioral and Pharmacotherapy Interventions <u>USPSTF Rating (Sept. 2015): A</u> • The USPSTF recommends that clinicians ask all <u>pregnant</u>	<ul> <li>Procedure Code(s): <u>Behavioral Interventions:</u></li> <li>99406, 99407</li> <li>99401, 99402, 99403, 99404</li> <li>Also see codes in the "Wellness Examinations" row above.</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code</li> </ul>	<ul> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> </ul>

Preventive Care Services

UnitedHealthcare Commercial Coverage Determination Guideline

Proprietary Information of UnitedHealthcare. Copyright 2017 United HealthCare Services, Inc.

#### Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
<ul> <li>women about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant women who use tobacco.</li> <li>The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and U.S. Food and Drug Administration (FDA)-approved pharmacotherapy for cessation to adults who use tobacco.</li> </ul>	requirements for preventive benefit to apply.	
Primary Care Interventions To Prevent Tobacco Use In Children And Adolescents <u>USPSTF Rating (Aug. 2013): B</u> The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents. <u>Bright Futures (April 2017):</u> Bright Futures recommends tobacco use assessments from age 11 years – 21 years.	<ul> <li>Procedure Code(s): Smoking and tobacco use cessation counseling visit:</li> <li>99406, 99407</li> <li>99401, 99402, 99403, 99404</li> <li>Also see codes in the "Wellness Examinations" row above.</li> <li>Diagnosis Code(s):</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>	<ul> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>
Screening for Visual Impairment in Children USPSTF Rating (Jan. 2011): B The USPSTF recommends vision screening for all children at least once between the ages of 3 and 5 years, to detect the presence of amblyopia or its risk factors.	<ul> <li>Procedure Code(s):</li> <li>99173, 99174, 99177</li> <li>Diagnosis Code(s):</li> <li>See Preventive Benefit Instructions column.</li> </ul>	<ul> <li>Age Limit (99173, 99174 and 99177):</li> <li>Less than age 6 years (ends on 6<sup>th</sup> birthday).</li> <li>Code 99173:</li> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> <li>Code 99174 and 99177:</li> <li>See the Medical Policy titled <u>Omnibus Codes</u> for allowable diagnoses.</li> </ul>
Behavioral Counseling to Prevent Skin Cancer <u>USPSTF Rating (May 2012): B</u> The USPSTF recommends counseling children, adolescents, and young adults aged 10 to 24 years who have fair skin about minimizing their exposure to	Procedure Code(s): <ul> <li>n/a</li> </ul> Diagnosis Code(s): <ul> <li>n/a</li> </ul>	<ul> <li>This service is included in a preventive care wellness examination or focused E&amp;M visit.</li> </ul>

Preventive Care Services

Page 23 of 50 Effective 10/01/2017

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
ultraviolet radiation to reduce risk for skin cancer.		
Prevention of Falls in Community-Dwelling Older Adults USPSTF Rating (May 2012): B The USPSTF recommends exercise or physical therapy and vitamin D supplementation to prevent falls in community- dwelling adults aged 65 years or older who are at increased risk for falls.	<pre>Procedure Code(s):     • n/a Diagnosis Code(s):     • n/a</pre>	<ul> <li>This service is included in a preventive care wellness examination or focused E&amp;M visit.</li> </ul>
Screening for Intimate Partner Violence <u>USPSTF Rating (Jan. 2013): B</u> The USPSTF recommends that clinicians screen women of childbearing age for intimate partner violence, such as domestic violence, and provide or refer women who screen positive to intervention services.	Procedure Code(s): <ul> <li>n/a</li> </ul> Diagnosis Code(s): <ul> <li>n/a</li> </ul> Procedure Code(s):	This service is included in a preventive care wellness examination.
Screening for Lung Cancer with Low-Dose Computed Tomography USPSTF Rating (Dec. 2013): B The USPSTF recommends annual screening for lung cancer with low-dose computed tomography in adults ages 55 to 80 years who have a 30 pack-year smoking history and currently smoke or have quit within the past 15 years. Screening should be discontinued once a person has not smoked for 15 years or develops a health problem that substantially limits life expectancy or the ability or willingness to have curative lung surgery.	<ul> <li>Procedure Code(s):</li> <li>G0297</li> <li>Diagnosis Code(s):</li> <li>ICD-10: F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891</li> <li>Codes for Reporting Purposes:</li> <li>G9275, G9276, G9458, G9459, G9460</li> <li>Note: codes G9275, G9276, G9458, G9459, and G9460 are for reporting purposes only, if applicable. These codes are not separately reimbursable.</li> </ul>	<ul> <li>Requires one of the listed diagnosis codes in this row.</li> <li>Limitations: <ul> <li>Limited to one per year, and</li> </ul> </li> <li>All of the following criteria: <ul> <li>Age 55 to 80 years (ends on 81<sup>st</sup> birthday), and</li> <li>At least 30 pack-years* of smoking history, and</li> <li>Either a current smoker, or, have quit within the past 15 years.</li> </ul> </li> <li>Note: Prior authorization requirements may apply, depending on plan.</li> <li>*A pack-year is a way to measure the amount a person has smoked over a long period of time. It is calculated by multiplying the number of packs of cigarettes smoked per day by the number of years the person has smoked. For example, 1 pack year is equal to smoking 1 pack per day for 1 year, or 2 packs per day for half a year, and so on. Source: National Institutes of Health, National Cancer Institute Dictionary of Cancer Terms, pack year definition web page. <a href="http://www.cancer.gov/dictionary?CdrID=30.6510">http://www.cancer.gov/dictionary?CdrID=30.6510</a></li> </ul>

Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

Tor preventiv	e care medications refer to pharm	nacy plan aunimistrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Fluoride Application in Primary Care USPSTF Rating (May 2014): B Children From Birth Through Age 5 Years. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption. Bright Futures (March 2014): For those at high risk, consider application of fluoride varnish for caries prevention. Recommended at 6 months, 9 months, 12 months, 18 months, 24 months, 30 months, 3 years and 6 years.	<ul> <li>Procedure Code(s): Application of topical fluoride by physician or other qualified health care professional: • 99188</li> <li>Diagnosis Code(s): • Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>	<ul> <li>Age 0 – 6 years (ends on 7th birthday)</li> <li>Does not have diagnosis code requirements for preventive benefit to apply.</li> </ul>
Latent Tuberculosis Infection: Screening, Adults USPSTF Rating (Sept. 2016): B The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk. This recommendation applies to asymptomatic adults <b>18 years</b> and older at increased risk for tuberculosis.	<ul> <li>Procedure Code(s):</li> <li>Screening: <ul> <li>86480, 86481, 86580</li> <li>99211 (for followup visit to check skin results)</li> </ul> </li> <li>Blood draw: <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>ICD-10:</li> <li>R76.11, R76.12</li> <li>Z00.00, Z00.01, Z11.1, Z20.1</li> </ul> </li> <li>Note: For age 18 – 20 years (ends on 21<sup>st</sup> birthday): in addition to the codes in this row, the preventive benefit also applies to the ICD-10 diagnosis codes listed in the Bright Futures row below, "Tuberculosis TB Testing"</li> </ul>	<ul> <li>Screening: <ul> <li>Ages 18 years and up.</li> <li>CPT code 86480, 86481, and 86580 are payable as preventive with any of the Diagnosis Codes listed in this row.</li> <li>CPT code 99211 is only payable as preventive with diagnosis code R76.11, or R76.12</li> </ul> </li> <li>Blood draw: <ul> <li>Ages 18 years and up. Payable as preventive when billed with 86480 or 86481, AND one of the Diagnosis codes listed in this row.</li> </ul> </li> </ul>
<b>Preeclampsia Screening</b> USPSTF Rating (April 2017): B The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy.	Preeclampsia screening by blood pressure measurement is included in the code for a wellness examination visit. See the codes in the Wellness Examinations row above.	See the Wellness Examinations row above.

Preventive Care Services

Also see the Expanded Women's Preventive Health table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Bright Futures:		
<b>Anemia Screening in Children</b> (Bright Futures)	Procedure Code(s): Anemia Screening in Children: ○ 85014, 85018 Blood draw: ○ 36415, 36416 Diagnosis Code(s): • <u>ICD-10:</u> Z00.110, Z00.111, Z00.121, Z00.129, Z13.0	<ul> <li>Anemia Screening in Children:</li> <li>Ages prenatal to 21 (ends on 21st birthday). No frequency limit. CPT codes 85014 and 85018 payable as preventive with one of the Diagnosis Codes listed in this row.</li> <li>Blood draw:</li> <li>Ages prenatal to 21 (ends on 21<sup>st</sup> birthday) payable when billed with 85014 or 85018, <u>AND</u> with one of the Diagnosis Codes listed in this row.</li> </ul>
Hearing Tests (Bright Futures, April 2017) <u>Hearing tests:</u> recommended at ages: Newborn; Between 3-5 days to 2 months; 4 years; 5 years, 6 years; 8 years;10 years; Once between age 11 – 14 years; Once between age 15 – 17 years; Once between age 18 – 21 years. Also recommended for those that have a positive risk assessment. <u>Risk assessment:</u> recommended at ages: 4 mo, 6 mo, 9 mo, 12 mo, 15 mo, 18 mo, 24 mo, 30 mo, 3 years, 7 years, 9 years	Procedure Code(s): Hearing tests: 92551, 92552, 92553 Diagnosis Code(s): <u>ICD-10:</u> Z00.121, Z00.129, Z01.10 Note: A risk assessment is included in the code for a wellness examination visit. See the codes in the Wellness Examinations row above.	<ul> <li>Ages prenatal to 21 (ends on 21st birthday). Limit of once per year. Payable as preventive with one of the Diagnosis Codes listed in this row.</li> </ul>
<ul> <li>Formal Developmental / Autism Screening (Bright Futures)</li> <li>A formal, standardized developmental screen is recommended during the 9 month visit.</li> <li>A formal, standardized developmental screen is recommended during the 18 month visit, including a formal autism screen.</li> <li>A formal, standardized autism screen is recommended during the 24 month visit.</li> <li>A formal, standardized developmental screen is recommended during the 24 month visit.</li> <li>A formal, standardized developmental screen is recommended during the 30 month visit.</li> </ul>	Procedure Code(s): ○ 96110 Diagnosis Code(s): ○ <u>ICD-10:</u> Z00.121, Z00.129, Z13.4	<ul> <li>Ages prenatal to 2 years (ends on 3rd birthday). No frequency limit. Payable as preventive with one of the Diagnosis Codes listed in this row.</li> </ul>

Preventive Care Services

#### Also see the <u>Expanded Women's Preventive Health</u> table below. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications refer to pharmacy plan administrator.

For preventive care medications refer to pharmacy plan administrator.		
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Lead Screening (Bright Futures)	Procedure Code(s): Lead Screening: o 83655 Blood draw: o 36415, 36416	<ul> <li>Lead Screening:</li> <li>Ages prenatal to 21 (ends on 21st birthday). No frequency limit. Payable as preventive with one of the Diagnosis Codes listed in this row.</li> </ul>
	Diagnosis Code(s): ○ <u>ICD-10</u> : Z00.121,Z00.129, Z77.011	<ul> <li>Blood draw:</li> <li>Ages prenatal to 21 (ends on 21<sup>st</sup> birthday) payable when billed with 83655</li> <li><u>AND</u> one of the Diagnosis Codes listed in this row.</li> </ul>
<b>Tuberculosis - TB Testing</b> (Bright Futures) <i>For age 18 years and older,</i> <i>please also refer to the USPSTF</i> <i>row above</i> "Latent Tuberculosis Infection: Screening, Adults"	<ul> <li>Procedure Code(s): <ul> <li>86580</li> <li>99211(for followup visit to check skin results)</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li><u>ICD-10</u>: R76.11, R76.12, Z20.1, Z00.121, Z00.129, Z11.1</li> </ul> </li> <li>Note: For age 18 years and up: <ul> <li>in addition to the codes in this row, the preventive benefit also applies to <u>all</u> codes listed in the USPSTF row above, "Latent Tuberculosis Infection: Screening, Adults".</li> </ul></li></ul>	<ul> <li>Ages prenatal to 21(ends on 21<sup>st</sup> birthday). For age 18 years and older, please also refer to the USPSTF row above "Latent Tuberculosis Infection: Screening, Adults"</li> <li>No frequency limit.</li> <li>CPT code 86580 is payable as preventive with one of the Diagnosis Codes listed in this row.</li> <li>CPT code 99211 is only payable as preventive with diagnosis code ICD-10: R76.11, R76.12 and Z11.1</li> </ul>
Dyslipidemia Screening (Bright Futures April 2014) Screening Lab Work: Conduct if risk assessment is positive, or, at the following intervals: once between age 9 – 11 years; once between age 17 – 21 years. Risk assessment: Recommended at 24 mo, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, 16 years.	<ul> <li>Procedure Code(s): Dyslipidemia Screening Lab work: <ul> <li>80061, 82465, 83718, 83719, 83721, 84478</li> </ul> </li> <li>Blood draw: <ul> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li><u>ICD-10</u>: Z00.121, Z00.129, Z13.220</li> </ul> </li> <li>Note: A risk assessment is included in the code for a wellness examination visit. See the Wellness Examinations row above.</li> </ul>	<ul> <li>Dyslipidemia Screening Lab Work:</li> <li>Ages 24 months to 21 years (ends on 21<sup>st</sup> birthday).</li> <li>Payable as preventive with one of the Diagnosis Codes listed in this row.</li> <li>Blood draw:</li> <li>Ages 24 months to 21 years (ends on 21<sup>st</sup> birthday) payable when billed with one of the listed Dyslipidemia Screening Procedure Codes listed in this row, AND with one of the Diagnosis Codes listed in this row.</li> </ul>
Tobacco, Alcohol or Drug Use Assessment (Bright Futures April 2017): Bright Futures recommends tobacco, alcohol or drug use assessment from age 11 years – 21 years.	<ul> <li>See codes in the following rows above:</li> <li>"Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents" and</li> <li>"Screening and Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse"</li> </ul>	<ul> <li>See the rows above:</li> <li>"Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents" and</li> <li>"Screening and Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse"</li> </ul>

Preventive Care Services

i or preventiv	e care medications refer to pharm	nacy plan administrator.
<b>Service:</b> A date in this column is when the rating was released, not when the benefit is effective.	Code(s):	Preventive Benefit Instructions:
Psychosocial / Behavioral Assessment Bright Futures (April 2017): Bright Futures recommends physicians conduct psychosocial / behavioral assessment at each of the recommended visits between newborn – 21 years.	An assessment is included in the code for a wellness examination visit. See the codes in the Wellness Examinations row above.	See the Wellness Examinations row above.
Depression Screening (Bright Futures, April 2017) Bright Futures recommends depression screening at each of the recommended visits between age 12 years – 21 years.	See the codes in the "Depression in Children and Adolescents (Screening) row above.	See the "Depression in Children and Adolescents (Screening)" row above.
Sexually Transmitted Infections (STI) Bright Futures (April 2017): Bright Futures recommends the following: STI Risk Assessment: Conduct risk assessment at each of the recommended visits between 11 years – 21 years. STI Lab Work: Conduct if risk assessment is positive	<ul> <li>See the codes in the following rows above:</li> <li>"Chlamydia Infection Screening"</li> <li>"Gonorrhea Screening"</li> </ul>	<ul> <li>See the following rows above:</li> <li>"Chlamydia Infection Screening"</li> <li>"Gonorrhea Screening"</li> </ul>
HIV Screening Bright Futures (April 2017) <u>HIV Risk Assessment:</u> Conduct risk assessment at age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years and 21 years. <u>HIV Screening Lab Work</u> : Conduct once between age 15 – 18 years. Also recommended anytime between ages 11 – 14 years, and 19 – 21 years when a risk assessment is positive.	See the codes in the "HIV – Human Immunodeficiency Virus – Screening for Adolescents and Adults" row above.	See the row above "HIV – Human Immunodeficiency Virus – Screening for Adolescents and Adults"

## **PREVENTIVE IMMUNIZATIONS**

An immunization that does not fall under one of the exclusions in the Certificate of Coverage is considered covered after the following conditions are satisfied: (1) FDA approval; (2) explicit ACIP recommendation published in the Morbidity & Mortality Weekly Report (MMWR) of the Centers for Disease Control and Prevention (CDC). Implementation will typically occur within 60 days after publication in the MMWR. In the case of a public health emergency (as defined by the Centers for Disease Control or state or local public health departments) UnitedHealthcare may choose to apply preventive benefits to a new vaccine if the vaccine has FDA approval, even if an ACIP recommendation has not been announced.

## NOTE:

- **Trade Name(s)** <u>column</u>: brand names/trade names are included, when available, as examples for convenience only. Coverage pursuant to this Coverage Determination Guideline is based solely on the procedure codes.
- Age Group <u>column</u>: This column is provided for informational use only. For purposes of this document: Adult means age 18 years and up; Pediatric means age 0-18 years.
- **Benefit Limits** <u>column</u>: Benefit Limits in **bold text** are from FDA labeling and ACIP recommendations. Codes that indicate "For applicable age see code description" are limited to the age(s) listed in the code description.

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
Immunization Administration Preventive when included as part of a preventive immunization.	90460	Immunization administration through 18 years of age via any route of administration, with counseling by physician or other qualified health care professional; first or only component of each vaccine or toxoid administered	n/a	Pediatric	For applicable age see code description.	
	90461	Immunization administration <b>through 18 years of age</b> via any route of administration, with counseling by physician or other qualified health care professional; each additional vaccine or toxoid component administered (List separately in addition to code for primary procedure)	n/a	Pediatric	For applicable age see code description.	
	90471	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); one vaccine (single or combination vaccine/toxoid)	n/a	Both	-	
	90472	Immunization administration (includes percutaneous, intradermal, subcutaneous, or intramuscular injections); each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure)	n/a	Both	-	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
	90473	Immunization administration by intranasal or oral route; one vaccine (single or combination vaccine/toxoid)	n/a	Both	-	
	90474	Immunization administration by intranasal or oral route; each additional vaccine (single or combination vaccine/toxoid) (List separately in addition to code for primary procedure)	n/a	Both	-	
	G0008	Administration of influenza virus vaccine	n/a	Both	-	
	G0009	Administration of pneumococcal vaccine	n/a	Both	-	
	G0010	Administration of hepatitis B vaccine	n/a	Both	-	
	0771 (revenue code)	Vaccine administration	n/a	Both	-	
Meningococcal (MenB-4C; MenB-Fhbp; Hib-MenCY; MPSV4;	90620	Meningococcal recombinant protein and outer membrane vesicle vaccine, serogroup B (MenB-4C), 2 dose schedule, for intramuscular use	• Bexsero®	Both	<u>Benefit Limit:</u> Age 10 and up.	
MCV4; MenACWY)	90621	Meningococcal recombinant lipoprotein vaccine, serogroup B (MenB-FHbp), 2 or 3 dose schedule, for intramuscular use	• Trumenba <sup>®</sup>	Both	<u>Benefit Limit:</u> Age 10 and up.	
	90644	Meningococcal conjugate vaccine, serogroups C & Y and Haemophilus influenzae b vaccine (Hib-MenCY), 4 dose schedule, when administered to children 2-15 months of age, for intramuscular use	• MenHibrix <sup>®</sup>	Pediatric	For applicable age see code description.	
	90733	Meningococcal polysaccharide vaccine, serogroups A, C, Y, W-135, quadrivalent (MPSV4) for subcutaneous use	• Menomune <sup>®</sup>	Both	-	
	90734	Meningococcal conjugate vaccine, serogroups A, C, Y and W-135, quadrivalent (MCV4 or MenACWY), for intramuscular use	<ul> <li>Menactra<sup>®</sup></li> <li>Menveo<sup>®</sup></li> </ul>	Both	-	
Hepatitis A	90632	Hepatitis A vaccine (HepA), adult dosage, for intramuscular use	• Havrix <sup>®</sup> • VAQTA <sup>®</sup>	Adult	For applicable age see code description.	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.					
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)
	90633	Hepatitis A vaccine (HepA), pediatric/ adolescent dosage-2 dose schedule, for intramuscular use	<ul> <li>Havrix<sup>®</sup></li> <li>VAQTA<sup>®</sup></li> </ul>	Pediatric	For applicable age see code description.
	90634	Hepatitis A vaccine (HepA), pediatric/adolescent dosage-3 dose schedule, for intramuscular use	• Havrix <sup>®</sup>	Pediatric	For applicable age see code description.
	90636	Hepatitis A and hepatitis B vaccine (HepA-HepB <b>), adult dosage,</b> for intramuscular use	• Twinrix <sup>®</sup>	Adult	For applicable age see code description.
Haemophilus influenza b (Hib):	90647	Haemophilus influenzae b vaccine (Hib), PRP-OMP conjugate, 3 dose schedule, for intramuscular use	• PedvaxHIB <sup>®</sup>	Both	-
	90648	Haemophilus influenzae b vaccine (Hib), PRP-T conjugate, 4 dose schedule, for intramuscular use	<ul> <li>ActHIB<sup>®</sup></li> <li>Hiberix<sup>®</sup></li> </ul>	Both	-
Human Papilloma Virus (HPV)	90649	Human Papilloma virus vaccine, types 6, 11, 16, 18, quadrivalent (HPV4), 3 dose schedule, for intramuscular use	• Gardasil4 <sup>®</sup>	Both	<u>Benefit Limit:</u> Ages 9-26yrs. Ends on 27 <sup>th</sup> birthday.
	90650	Human Papilloma virus vaccine, types 16, 18, bivalent (HPV2), 3 dose schedule, for intramuscular use	• -	Both	<u>Benefit Limit:</u> Ages 9-26yrs. Ends on 27 <sup>th</sup> birthday.
	90651	Human Papillomavirus vaccine types 6, 11, 16, 18, 31, 33, 45, 52, 58, nonavalent (9vHPV), 2 or 3 dose schedule, for intramuscular use	• Gardasil9 <sup>®</sup>	Both	<u>Benefit Limit:</u> Ages 9-26yrs. Ends on 27 <sup>th</sup> birthday.
Seasonal Influenza (`flu')	90630	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, for intradermal use	<ul> <li>Fluzone<sup>®</sup> Intradermal Quadrivalent</li> </ul>	Both	-
<u>Note</u> : Additional new seasonal flu immunization	90653	Influenza vaccine, inactivated (IIV), subunit, adjuvanted, for intramuscular use	-	Both	-
codes that are recently FDA- approved, but are not listed here, may be eligible for preventive benefits as of the FDA approval date.	90654	Influenza virus vaccine, trivalent (IIV3), split virus, preservative-free, for intradermal use	<ul> <li>Fluzone<sup>®</sup> Intradermal Trivalent</li> </ul>	Adult	<u>Benefit Limit:</u> 18 years – 64 years. Ends on 65 <sup>th</sup> birthday.

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	<b>Trade</b> <b>Name(s)</b> (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
	90655	Influenza virus vaccine, trivalent (IIV3), split virus, preservative free, 0.25 mL dosage, for intramuscular use	<ul> <li>Fluzone<sup>®</sup> No Preservative Pediatric</li> </ul>	Pediatric	<u>Benefit Limit:</u> 6 – 35 months old.	
	90656	Influenza virus vaccine, trivalent (IIV3), split virus, preservative free, 0.5 mL dosage, for intramuscular use	<ul> <li>Afluria<sup>®</sup></li> <li>Fluzone<sup>®</sup> No preservative</li> <li>Fluvirin<sup>®</sup></li> <li>Fluarix<sup>®</sup></li> <li>Flulaval<sup>®</sup></li> </ul>	Both	Benefit Limit: 3 years and up.	
	90657	Influenza virus vaccine, trivalent (IIV3), split virus, 0.25 mL dosage, for intramuscular use	• Fluzone <sup>®</sup>	Pediatric	<u>Benefit Limit:</u> 6 – 35 months old.	
	90658	Influenza virus vaccine, trivalent (IIV3), split virus, 0.5 mL dosage, for intramuscular use	<ul> <li>Afluria<sup>®</sup></li> <li>Flulaval<sup>®</sup></li> <li>Fluvirin<sup>®</sup></li> <li>Fluzone<sup>®</sup></li> </ul>	Both	Benefit Limit: 3 years and up.	
	90660	Influenza virus vaccine, trivalent, live (LAIV3), for intranasal use	• Flumist <sup>®</sup>	Both	<u>Benefit Limit:</u> Ages 2 - 49 Years. Ends on 50 <sup>th</sup> birthday	
	90661	Influenza virus vaccine, trivalent (ccIIV3), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, for intramuscular use	<ul> <li>Flucelvax<sup>™</sup></li> </ul>	Adult	<u>Benefit Limit:</u> Ages 4 years and up	
	90662	Influenza virus vaccine (IIV), split virus, preservative free, enhanced immunogenicity via increased antigen content, for intramuscular use	<ul> <li>High Dose Fluzone<sup>®</sup></li> </ul>	Adult	<u>Benefit Limit:</u> Ages 65 years and up	
	90664	Influenza virus vaccine, live (LAIV), pandemic formulation, for intranasal use	• Flumist <sup>®</sup>	Both	<u>Benefit Limit:</u> Ages 2 - 49 Years. Ends on 50 <sup>th</sup> birthday.	
	90666	Influenza virus vaccine (IIV), pandemic formulation, split virus, preservative free, for intramuscular use	-	Both	-	
	90667	Influenza virus vaccine (IIV), pandemic formulation, split virus, adjuvanted, for intramuscular use	-	Both	-	
	90668	Influenza virus vaccine (IIV), pandemic formulation, split virus, for intramuscular use	-	Both	-	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
	90672	Influenza virus vaccine, quadrivalent, live (LAIV4), for intranasal use	• Flumist <sup>®</sup> (LAIV4)	Both	<u>Benefit Limit:</u> Ages 2 – 49 Years. Ends on 50 <sup>th</sup> birthday.	
	90673	Influenza virus vaccine, trivalent (RIV3), derived from recombinant DNA (RIV3), hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use	• Flublok <sup>®</sup>	Adult	<u>Benefit Limit:</u> Age 18 years and up.	
	90674	Influenza virus vaccine, quadrivalent (ccIIV4), derived from cell cultures, subunit, preservative and antibiotic free, 0.5 mL dosage, for intramuscular use	• FLUCELVAX <sup>®</sup> QUADRIVALENT	Both	<u>Benefit Limit:</u> Age 4 years and up.	
	90682	Influenza virus vaccine, quadrivalent (RIV4), derived from recombinant DNA, hemagglutinin (HA) protein only, preservative and antibiotic free, for intramuscular use	<ul> <li>Flublok Quadrivalent</li> </ul>	Adult	<u>Benefit Limit:</u> Age 18 years and up.	
	90685	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.25 mL, for intramuscular use	• Fluzone <sup>®</sup>	Pediatric	<u>Benefit Limit:</u> 6 – 35 months old.	
	90686	Influenza virus vaccine, quadrivalent (IIV4), split virus, preservative free, 0.5 mL dosage, for intramuscular use	<ul> <li>Fluarix<sup>®</sup></li> <li>FluLaval Quadrivalent <sup>®</sup></li> <li>Fluzone Quadrivalent <sup>®</sup></li> </ul>	Both	Benefit Limit: Ages 6 months and up.	
	90687	Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.25 mL dosage, for intramuscular use	<ul> <li>Fluzone Quadrivalent ®</li> </ul>	Pediatric	<u>Benefit Limit:</u> 6 – 35 months old.	
	90688	Influenza virus vaccine, quadrivalent (IIV4), split virus, 0.5 mL dosage, for intramuscular use	<ul> <li>FluLaval Quadrivalent         <ul> <li>Fluzone Quadrivalent             </li> </ul> </li> </ul>	Both	<u>Benefit Limit:</u> Ages 6 months and up.	
	Q2034	Influenza virus vaccine, split virus, for intramuscular use (Agriflu)	• Agriflu <sup>®</sup>	Adult	<u>Benefit Limit:</u> Ages 18 years and up	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
	Q2035	Influenza virus vaccine, split virus, when administered to individuals <b>3 years of age</b> <b>and older,</b> for intramuscular use (AFLURIA)	• Afluria <sup>®</sup>	Both	For applicable age see code description.	
	Q2036	Influenza virus vaccine, split virus, when administered to individuals <b>3 years of age</b> <b>and older,</b> for intramuscular use (FLULAVAL)	• Flulaval <sup>®</sup>	Both	For applicable age see code description.	
	Q2037	Influenza virus vaccine, split virus, when administered to individuals <b>3 years of age</b> <b>and older,</b> for intramuscular use (FLUVIRIN)	• Fluvirin <sup>®</sup>	Both	For applicable age see code description.	
	Q2038	Influenza virus vaccine, split virus, when administered to individuals <b>3 years of age</b> <b>and older</b> , for intramuscular use (Fluzone)	• Fluzone <sup>®</sup>	Both	For applicable age see code description.	
	Q2039	Influenza virus vaccine, not otherwise specified	-	Both	-	
Pneumococcal polysaccharide (PPSV23)	90732	Pneumococcal polysaccharide vaccine, 23-valent (PPSV23), adult or immunosuppressed patient dosage, when administered to <b>individuals</b> <b>2 years or older,</b> for subcutaneous or intramuscular use	• Pneumovax 23 <sup>®</sup>	Both	For applicable age see code description.	
Pneumococcal conjugate	90670	Pneumococcal conjugate vaccine, 13 valent (PCV13), for intramuscular use	<ul> <li>Prevnar 13<sup>®</sup> (PCV13)</li> </ul>	Both	-	
Rotavirus	90680	Rotavirus vaccine, pentavalent (RV5), 3 dose schedule, live, for oral use	• ROTATEQ <sup>®</sup>	Both	-	
	90681	Rotavirus vaccine, human, attenuated (RV1), 2 dose schedule, live, for oral use	• Rotarix <sup>®</sup>	Both	-	
Diphtheria, tetanus toxoids, acellular pertussis and polio inactive (DTap-IPV)	90696	Diphtheria, tetanus toxoids, acellular pertussis vaccine and inactivated poliovirus vaccine (DTaP-IPV), when administered to <b>children 4</b> <b>through 6 years of age,</b> for intramuscular use	<ul> <li>Kinrix<sup>®</sup></li> <li>Quadracel<sup>®</sup></li> </ul>	Pediatric	For applicable age see code description.	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.						
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)	
Diphtheria, tetanus toxoids, acellular pertussis, haemophilus influenza B, and polio inactive (DTap-IPV/Hib)	90698	Diphtheria, tetanus toxoids, acellular pertussis vaccine, Haemophilus influenzae type b, and inactivated poliovirus vaccine, (DTaP-IPV/Hib), for intramuscular use	• Pentacel <sup>®</sup>	Both	-	
Diphtheria, tetanus, acellular pertussis (DTap)	90700	Diphtheria, tetanus toxoids, and acellular pertussis vaccine (DTaP), when administered to <b>individuals</b> <b>younger than 7 years</b> , for intramuscular use	<ul> <li>Daptacel<sup>®</sup></li> <li>Infanrix<sup>®</sup></li> </ul>	Pediatric	For applicable age see code description.	
Diphtheria and tetanus (DT)	90702	Diphtheria and tetanus toxoids adsorbed (DT) when administered to individuals <b>younger than 7 years,</b> for intramuscular use	-	Pediatric	For applicable age see code description.	
Measles, Mumps, Rubella (MMR)	90707	Measles, mumps and rubella virus vaccine (MMR), live, for subcutaneous use	• MMR II <sup>®</sup>	Both	-	
	90710	Measles, mumps, rubella, and varicella vaccine (MMRV), live, for subcutaneous use	• ProQuad <sup>®</sup>	Both	-	
Polio (IPV)	90713	Poliovirus vaccine, inactivated (IPV), for subcutaneous or intramuscular use	<ul> <li>Ipol<sup>®</sup></li> </ul>	Both	-	
Tetanus and diphtheria (Td)	90714	Tetanus and diphtheria toxoids adsorbed (Td), preservative free, when administered to <b>individuals</b> <b>7 years or older,</b> for intramuscular use	<ul> <li>Tenivac<sup>®</sup></li> <li>Decavac<sup>®</sup></li> </ul>	Both	For applicable age see code description.	
Tetanus, diphtheria toxoids and acellular pertussis (Tdap)	90715	Tetanus, diphtheria toxoids and acellular pertussis vaccine (Tdap), when administered <b>to individuals</b> <b>7 years or older</b> , for intramuscular use	<ul> <li>Adacel<sup>®</sup></li> <li>Boostrix<sup>®</sup></li> </ul>	Both	For applicable age see code description.	
Varicella (VAR) (`chicken pox')	90716	Varicella virus vaccine (VAR), live, for subcutaneous use	• Varivax <sup>®</sup>	Both	-	
Diptheria, tetanus and acellular pertussis, hep B, and polio inactive (DTaP-HepB-IPV)	90723	Diphtheria, tetanus toxoids, acellular pertussis vaccine, hepatitis B, and inactivated poliovirus vaccine (DTaP- HepB-IPV), for intramuscular use	• PEDIARIX <sup>®</sup>	Both	<u>Benefit Limit:</u> Ages 0-6yrs. Ends on 7 <sup>th</sup> birthday.	

<b>PREVENTIVE IMMUNIZATIONS</b> These codes do <b>not</b> have a diagnosis code requirement for preventive benefits to apply.							
Category:	Code(s):	Description:	Trade Name(s) (See NOTE above)	Age Group: • Adult (18yr.+) • Pediatric (0-18yr), • Both	Benefit Limits: Age / Other (See NOTE above)		
Zoster / Shingles (HZV)	90736	Zoster (shingles) vaccine (HZV), live, for subcutaneous injection	<ul> <li>Zostavax</li> </ul>	Adult	<u>Benefit Limit:</u> Age 60 years and up.		
Hepatitis B	90740	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 3 dose schedule, for intramuscular use	• Recombivax HB <sup>®</sup>	Both	-		
	90743	Hepatitis B vaccine (HepB), adolescent, 2 dose schedule, for intramuscular use	<ul> <li>Recombivax HB<sup>®</sup></li> </ul>	Pediatric (adolesce nt only)	For applicable age see code description.		
	90744	Hepatitis B vaccine (HepB), pediatric/adolescent dosage, 3 dose schedule, for intramuscular use	<ul> <li>Recombivax HB<sup>®</sup></li> <li>Energix-B<sup>®</sup></li> </ul>	Pediatric	For applicable age see code description.		
	90746	Hepatitis B vaccine (HepB), adult dosage, 3 dose schedule, for intramuscular use	<ul> <li>Recombivax HB<sup>®</sup></li> <li>Energix-B<sup>®</sup></li> </ul>	Adult	For applicable age see code description.		
	90747	Hepatitis B vaccine (HepB), dialysis or immunosuppressed patient dosage, 4 dose schedule, for intramuscular use	• Energix-B <sup>®</sup>	Both	-		
	90748	Hepatitis B and Haemophilus influenza b vaccine (Hib- HepB), for intramuscular use	-	Both	-		

## Pregnancy Diagnosis Code List:

The following Pregnancy Diagnosis Codes are required where indicated in the Preventive Care Services table (above), or in the Expanded Women's Preventive Health table (below):

## ICD-10 Codes:

000.00, 000.01, 000.101, 000.102, 000.109, 000.111, 000.112, 000.119, 000.201, 000.202, 000.209, 000.211, 000.212, 000.219, 000.80, 000.81, 000.90, 000.91, 009.A0, 009.A1, 009.A2, 009.A3, 001.0, 001.1, 001.9, 002.0, 002.1, 002.81, 002.89, 002.9, 003.0, 003.1, 003.2, 003.30, 003.31, 003.32, 003.33, 003.34, 003.35, 003.36, 003.37, 003.38, 003.39, 003.4, 003.5, 003.6, 003.7, 003.80, 003.81, 003.82, 003.83, 003.84, 003.85, 003.86, 003.87, 003.88, 003.89, 003.9, 004.5, 004.6, 004.7, 004.80, 004.81, 004.82, 004.83, 004.84, 004.85, 004.86, 004.87, 004.88, 004.89, 007.0, 007.1, 007.2, 007.30, 007.31, 007.32, 007.33, 007.34, 007.35, 007.36, 007.37, 007.38, 007.39, 007.4, 008.0, 008.1, 008.2, 008.3, 008.4, 008.5, 008.6, 008.7, 008.81, 008.82, 008.83, 008.89, 008.9, 009.00, 009.01, 009.02, 009.03, 009.10, 009.11, 009.12, 009.13, 009.211, 009.212, 009.213, 009.219, 009.291, 009.292, 009.293, 009.299, 009.30, 009.31, 009.32, 009.33, 009.40, 009.41, 009.42, 009.43, 009.511, 009.512, 009.513, 009.519, 009.521, 009.522, 009.523, 009.529, 009.611, 009.612, 009.613, 009.619, 009.621, 009.622, 009.623, 009.629, 009.70, 009.71, 009.72, 009.73, 009.811, 009.812, 009.813, 009.819, 009.821, 009.822, 009.823, 009.829, 009.891, 009.892, 009.893, 009.899, 009.90, 009.91, 009.92, 009.93, 010.011, 010.012, 010.013, 010.019, 010.02, 010.03, 010.111, 010.112, 010.113, 010.119, 010.12, 010.13, 010.211, 010.212, 010.213, 010.219, 010.22, 010.23, 010.311, 010.312, 010.313, 010.319, 010.32, 010.33, 010.411, 010.412, 010.413, 010.419, 010.42, 010.43, 010.911, 010.912, 010.913, 010.919, 010.92, 010.93, 011.1, 011.2, 011.3, 011.4, 011.5, 011.9, 012.00, 012.01, 012.02, 012.03, 012.10, 012.11, 012.12, 012.13, 012.20, 012.21, 012.22, 012.23, 012.04, 012.05, 012.14, 012.15, 012.24, 012.25, 013.1,

#### Pregnancy Diagnosis Code List:

The following Pregnancy Diagnosis Codes are required where indicated in the Preventive Care Services table (above), or in the Expanded Women's Preventive Health table (below):

013.2, 013.3, 013.4, 013.5, 013.9, 014.00, 014.02, 014.03, 014.04, 014.05, 014.10, 014.12, 014.13, 014.14, 014.15, 014.20, 014.22, 014.23, 014.24, 014.25, 014.90, 014.92, 014.93, 014.94, 014.95, 015.00, 015.02, 015.03, 015.1, 015.2, 015.9, 016.1, 016.2, 016.3, 016.4, 016.5, 016.9, 020.0, 020.8, 020.9, 021.0, 021.1, 021.2, 021.8, 021.9, 022.00, 022.01, 022.02, 022.03, 022.10, 022.11, 022.12, 022.13, 022.20, 022.21, 022.22, 022.23, 022.30, 022.31, 022.32, 022.33, 022.40, 022.41, 022.42, 022.43, 022.50, 022.51, 022.52, 022.53, 022.8X1, 022.8X2, 022.8X3, 022.8X9, 022.90, 022.91, 022.92, 022.93, 023.00, 023.01, 023.02, 023.03, 023.10, 023.11, 023.12, 023.13, 023.20, 023.21, 023.22, 023.23, 023.30, 023.31, 023.32, 023.33, 023.40, 023.41, 023.42, 023.43, 023.511, 023.512, 023.513, 023.519, 023.521, 023.522, 023.523, 023.529, 023.591, 023.592, 023.593, 023.599, 023.90, 023.91, 023.92, 023.93, 024.011, 024.012, 024.013, 024.019, 024.02, 024.03, 024.111, 024.112, 024.113, 024.119, 024.12, 024.13, 024.311, 024.312, 024.313, 024.319, 024.32, 024.33, 024.410, 024.414, 024.415, 024.419, 024.420, 024.424, 024.425, 024.429, 024.430, 024.434, 024.435, 024.439, 024.811, 024.812, 024.813, 024.819, 024.82, 024.83, 024.911, 024.912, 024.913, 024.919, 024.92, 024.93, 025.10, 025.11, 025.12, 025.13, 025.2, 025.3, 026.00, 026.01, 026.02, 026.03, 026.10, 026.11, 026.12, 026.13, 026.20, 026.21, 026.22, 026.23, 026.30, 026.31, 026.32, 026.33, 026.40, 026.41, 026.42, 026.43, 026.50, 026.51, 026.52, 026.53, 026.611, 026.612, 026.613, 026.619, 026.62, 026.63, 026.711, 026.712, 026.713, 026.719, 026.72, 026.73, 026.811, 026.812, 026.813, 026.819, 026.821, 026.822, 026.823, 026.829, 026.831, 026.832, 026.833, 026.839, 026.841, 026.842, 026.843, 026.849, 026.851, 026.852, 026.853, 026.859, 026.86, 026.872, 026.873, 026.879, 026.891, 026.892, 026.893, 026.899, 026.90, 026.91, 026.92, 026.93, 028.0, 028.1, 028.2, 028.3, 028.4, 028.5, 028.8, 028.9, 029.011, 029.012, 029.013, 029.019, 029.021, 029.022, 029.023, 029.029, 029.091, 029.092, 029.093, 029.099, 029.111, 029.112, 029.113, 029.119, 029.121, 029.122, 029.123, 029.129, 029.191, 029.192, 029.193, 029.199, 029.211, 029.212, 029.213, 029.219, 029.291, 029.292, 029.293, 029.299, 029.3X1, 029.3X2, 029.3X3, 029.3X9, 029.40, 029.41, 029.42, 029.43, 029.5X1, 029.5X2, 029.5X3, 029.5X9, 029.60, 029.61, 029.62, 029.63, 029.8X1, 029.8X2, 029.8X3, 029.8X9, 029.90, 029.91, 029.92, 029.93, 030.001, 030.002, 030.003, 030.009, 030.011, 030.012, 030.013, 030.019, 030.021, 030.022, 030.023, 030.029, 030.031, 030.032, 030.033, 030.039, 030.041, 030.042, 030.043, 030.049, 030.091, 030.092, 030.093, 030.099, 030.101, 030.102, 030.103, 030.109, 030.111, 030.112, 030.113, 030.119, 030.121, 030.122, 030.123, 030.129, 030.191, 030.192, 030.193, 030.199, 030.201, 030.202, 030.203, 030.209, 030.211, 030.212, 030.213, 030.219, 030.221, 030.222, 030.223, 030.229, 030.291, 030.292, 030.293, 030.299, 030.801, 030.802, 030.803, 030.809, 030.811, 030.812, 030.813, 030.819, 030.821, 030.822, 030.823, 030.829, 030.891, 030.892, 030.893, 030.899, 030.90, 030.91, 030.92, 030.93, 031.00X0, 031.00X1, 031.00X2, 031.00X3, 031.00X4, 031.00X5, 031.00X9, 031.01X0, 031.01X1, 031.01X2, 031.01X3, 031.01X4, 031.01X5, 031.01X9, 031.02X0, 031.02X1, 031.02X2, 031.02X3, 031.02X4, 031.02X5, 031.02X9, 031.03X0, 031.03X1, 031.03X2, 031.03X3, 031.03X4, 031.03X5, 031.03X9, O31.10X0, O31.10X1, O31.10X2, O31.10X3, O31.10X4, O31.10X5, O31.10X9, O31.11X0, O31.11X1, O31.11X2, 031.11X3, 031.11X4, 031.11X5, 031.11X9, 031.12X0, 031.12X1, 031.12X2, 031.12X3, 031.12X4, 031.12X5, 031.12X9, 031.13X0, 031.13X1, 031.13X2, 031.13X3, 031.13X4, 031.13X5, 031.13X9, 031.20X0, 031.20X1, 031.20X2, 031.20X3, 031.20X4, 031.20X5, 031.20X9, 031.21X0, 031.21X1, 031.21X2, 031.21X3, 031.21X4, 031.21X5, 031.21X9, 031.22X0, 031.22X1, 031.22X2, 031.22X3, 031.22X4, 031.22X5, 031.22X9, 031.23X0, 031.23X1, 031.23X2, 031.23X3, 031.23X4, 031.23X5, 031.23X9, 031.30X0, 031.30X1, 031.30X2, 031.30X3, 031.30X4, 031.30X5, 031.30X9, 031.31X0, 031.31X1, 031.31X2, 031.31X3, 031.31X4, 031.31X5, 031.31X9, 031.32X0, 031.32X1, 031.32X2, 031.32X3, 031.32X4, 031.32X5, 031.32X9, 031.33X0, 031.33X1, 031.33X2, 031.33X3, 031.33X4, 031.33X5, 031.33X9, 031.8X10, 031.8X11, 031.8X12, 031.8X13, 031.8X14, 031.8X15, 031.8X19, 031.8X20, 031.8X21, 031.8X22, 031.8X23, 031.8X24, 031.8X25, 031.8X29, 031.8X30, 031.8X31, 031.8X32, 031.8X33, 031.8X34, 031.8X35, 031.8X39, 031.8X90, 031.8X91, 031.8X92, 031.8X93, 031.8X94, O31.8X95, O31.8X99, O32.0XX0, O32.0XX1, O32.0XX2, O32.0XX3, O32.0XX4, O32.0XX5, O32.0XX9, O32.1XX0, 032.1XX1, 032.1XX2, 032.1XX3, 032.1XX4, 032.1XX5, 032.1XX9, 032.2XX0, 032.2XX1, 032.2XX2, 032.2XX3, 032.2XX4, 032.2XX5, 032.2XX9, 032.3XX0, 032.3XX1, 032.3XX2, 032.3XX3, 032.3XX4, 032.3XX5, 032.3XX9, 032.4XX0, 032.4XX1, 032.4XX2, 032.4XX3, 032.4XX4, 032.4XX5, 032.4XX9, 032.6XX0, 032.6XX1, 032.6XX2, 032.6XX3, 032.6XX4, 032.6XX5, 032.6XX9, 032.8XX0, 032.8XX1, 032.8XX2, 032.8XX3, 032.8XX4, 032.8XX5, 032.8XX9, 032.9XX0, 032.9XX1, 032.9XX2, 032.9XX3, 032.9XX4, 032.9XX5, 032.9XX9, 033.0, 033.1, 033.2, 033.3XX0, 033.3XX1, 033.3XX2, 033.3XX3, 033.3XX4, 033.3XX5, 033.3XX9, 033.4XX0, 033.4XX1, 033.4XX2, 033.4XX3, 033.4XX4, 033.4XX5, 033.4XX9, 033.5XX0, 033.5XX1, 033.5XX2, 033.5XX3, 033.5XX4, 033.5XX5, O33.5XX9, O33.6XX0, O33.6XX1, O33.6XX2, O33.6XX3, O33.6XX4, O33.6XX5, O33.6XX9, O33.7XX0, O33.7XX1, 033.7XX2, 033.7XX3, 033.7XX4, 033.7XX5, 033.7XX9033.8, 033.9, 034.00, 034.01, 034.02, 034.03, 034.10, 034.11, 034.12, 034.13, 034.211, 034.212, 034.219, 034.29, 034.30, 034.31, 034.32, 034.33, 034.40, 034.41, 034.42, 034.43, 034.511, 034.512, 034.513, 034.519, 034.521, 034.522, 034.523, 034.529, 034.531, 034.532, 034.533, 034.539, 034.591, 034.592, 034.593, 034.599, 034.60, 034.61, 034.62, 034.63, 034.70, 034.71, 034.72, 034.73, 034.80, 034.81, 034.82, 034.83, 034.90, 034.91, 034.92, 034.93, 035.0XX0, 035.0XX1, 035.0XX2, 035.0XX3, 035.0XX4, 035.0XX5, 035.0XX9, 035.1XX0, 035.1XX1, 035.1XX2, 035.1XX3, 035.1XX4,

# Pregnancy Diagnosis Code List:

The following Pregnancy Diagnosis Codes are required where indicated in the Preventive Care Services table (above), or in the Expanded Women's Preventive Health table (below):

or in the Expanded women's Preventive Health table (below).
O35.1XX5, O35.1XX9, O35.2XX0, O35.2XX1, O35.2XX2, O35.2XX3, O35.2XX4, O35.2XX5, O35.2XX9, O35.3XX0,
O35.3XX1, O35.3XX2, O35.3XX3, O35.3XX4, O35.3XX5, O35.3XX9, O35.4XX0, O35.4XX1, O35.4XX2, O35.4XX3,
O35.4XX4, O35.4XX5, O35.4XX9, O35.5XX0, O35.5XX1, O35.5XX2, O35.5XX3, O35.5XX4, O35.5XX5, O35.5XX9,
035.6XX0, 035.6XX1, 035.6XX2, 035.6XX3, 035.6XX4, 035.6XX5, 035.6XX9, 035.7XX0, 035.7XX1, 035.7XX2,
O35.7XX3, O35.7XX4, O35.7XX5, O35.7XX9, O35.8XX0, O35.8XX1, O35.8XX2, O35.8XX3, O35.8XX4, O35.8XX5,
O35.8XX9, O35.9XX0, O35.9XX1, O35.9XX2, O35.9XX3, O35.9XX4, O35.9XX5, O35.9XX9, O36.0110, O36.0111,
036.0112, 036.0113, 036.0114, 036.0115, 036.0119, 036.0120, 036.0121, 036.0122, 036.0123, 036.0124,
036.0125, 036.0129, 036.0130, 036.0131, 036.0132, 036.0133, 036.0134, 036.0135, 036.0139, 036.0190,
036.0191, 036.0192, 036.0193, 036.0194, 036.0195, 036.0199, 036.0910, 036.0911, 036.0912, 036.0913,
O36.0914, O36.0915, O36.0919, O36.0920, O36.0921, O36.0922, O36.0923, O36.0924, O36.0925, O36.0929,
036.0930, 036.0931, 036.0932, 036.0933, 036.0934, 036.0935, 036.0939, 036.0990, 036.0991, 036.0992,
036.0993, 036.0994, 036.0995, 036.0999, 036.1110, 036.1111, 036.1112, 036.1113, 036.1114, 036.1115,
036.1119, 036.1120, 036.1121, 036.1122, 036.1123, 036.1124, 036.1125, 036.1129, 036.1130, 036.1131,
036.1132, 036.1133, 036.1134, 036.1135, 036.1139, 036.1190, 036.1191, 036.1192, 036.1193, 036.1194,
036.1195, 036.1199, 036.1910, 036.1911, 036.1912, 036.1913, 036.1914, 036.1915, 036.1919, 036.1920,
036.1921, 036.1922, 036.1923, 036.1924, 036.1925, 036.1929, 036.1930, 036.1931, 036.1932, 036.1933,
036.1934, 036.1935, 036.1939, 036.1990, 036.1991, 036.1992, 036.1993, 036.1994, 036.1995, 036.1999,
O36.20X0, O36.20X1, O36.20X2, O36.20X3, O36.20X4, O36.20X5, O36.20X9, O36.21X0, O36.21X1, O36.21X2,
036.21X3, 036.21X4, 036.21X5, 036.21X9, 036.22X0, 036.22X1, 036.22X2, 036.22X3, 036.22X4, 036.22X5,
O36.22X9, O36.23X0, O36.23X1, O36.23X2, O36.23X3, O36.23X4, O36.23X5, O36.23X9, O36.4XX0, O36.4XX1,
O36.4XX2, O36.4XX3, O36.4XX4, O36.4XX5, O36.4XX9, O36.5110, O36.5111, O36.5112, O36.5113, O36.5114,
036.5115, 036.5119, 036.5120, 036.5121, 036.5122, 036.5123, 036.5124, 036.5125, 036.5129, 036.5130,
036.5131, 036.5132, 036.5133, 036.5134, 036.5135, 036.5139, 036.5190, 036.5191, 036.5192, 036.5193,
036.5194, 036.5195, 036.5199, 036.5910, 036.5911, 036.5912, 036.5913, 036.5914, 036.5915, 036.5919,
036.5920, 036.5921, 036.5922, 036.5923, 036.5924, 036.5925, 036.5929, 036.5930, 036.5931, 036.5932,
036.5933, 036.5934, 036.5935, 036.5939, 036.5990, 036.5991, 036.5992, 036.5993, 036.5994, 036.5995,
O36.5999, O36.60X0, O36.60X1, O36.60X2, O36.60X3, O36.60X4, O36.60X5, O36.60X9, O36.61X0, O36.61X1,
O36.61X2, O36.61X3, O36.61X4, O36.61X5, O36.61X9, O36.62X0, O36.62X1, O36.62X2, O36.62X3, O36.62X4,
O36.62X5, O36.62X9, O36.63X0, O36.63X1, O36.63X2, O36.63X3, O36.63X4, O36.63X5, O36.63X9, O36.70X0,
O36.70X1, O36.70X2, O36.70X3, O36.70X4, O36.70X5, O36.70X9, O36.71X0, O36.71X1, O36.71X2, O36.71X3,
036.71X4, 036.71X5, 036.71X9, 036.72X0, 036.72X1, 036.72X2, 036.72X3, 036.72X4, 036.72X5, 036.72X9,
O36.73X0, O36.73X1, O36.73X2, O36.73X3, O36.73X4, O36.73X5, O36.73X9, O36.80X0, O36.80X1, O36.80X2,
O36.80X3, O36.80X4, O36.80X5, O36.80X9, O36.8120, O36.8121, O36.8122, O36.8123, O36.8124, O36.8125,
036.8129, 036.8130, 036.8131, 036.8132, 036.8133, 036.8134, 036.8135, 036.8139, 036.8190, 036.8191,
036.8192, 036.8193, 036.8194, 036.8195, 036.8199, 036.8210, 036.8211, 036.8212, 036.8213, 036.8214,
036.8215, 036.8219, 036.8220, 036.8221, 036.8222, 036.8223, 036.8224, 036.8225, 036.8229, 036.8230,
036.8231, 036.8232, 036.8233, 036.8234, 036.8235, 036.8239, 036.8290, 036.8291, 036.8292, 036.8293,
036.8294, 036.8295, 036.8299, 036.8310, 036.8311, 036.8312, 036.8313, 036.8314, 036.8315, 036.8319,
036.8320, 036.8321, 036.8322, 036.8323, 036.8324, 036.8325, 036.8329, 036.8330, 036.8331, 036.8332,
036.8333, 036.8334, 036.8335, 036.8339, 036.8390, 036.8391, 036.8392, 036.8393, 036.8394, 036.8395,
036.8399, 036.8910, 036.8911, 036.8912, 036.8913, 036.8914, 036.8915, 036.8919, 036.8920, 036.8921,
036.8922, 036.8923, 036.8924, 036.8925, 036.8929, 036.8930, 036.8931, 036.8932, 036.8933, 036.8934,
O36.8935, O36.8939, O36.8990, O36.8991, O36.8992, O36.8993, O36.8994, O36.8995, O36.8999, O36.90X0,
O36.90X1, O36.90X2, O36.90X3, O36.90X4, O36.90X5, O36.90X9, O36.91X0, O36.91X1, O36.91X2, O36.91X3,
036.91X4, 036.91X5, 036.91X9, 036.92X0, 036.92X1, 036.92X2, 036.92X3, 036.92X4, 036.92X5, 036.92X9,
O36.93X0, O36.93X1, O36.93X2, O36.93X3, O36.93X4, O36.93X5, O36.93X9, O40.1XX0, O40.1XX1, O40.1XX2,
O40.1XX3, O40.1XX4, O40.1XX5, O40.1XX9, O40.2XX0, O40.2XX1, O40.2XX2, O40.2XX3, O40.2XX4, O40.2XX5,
O40.2XX9, O40.3XX0, O40.3XX1, O40.3XX2, O40.3XX3, O40.3XX4, O40.3XX5, O40.3XX9, O40.9XX0, O40.9XX1,
O40.9XX2, O40.9XX3, O40.9XX4, O40.9XX5, O40.9XX9, O41.00X0, O41.00X1, O41.00X2, O41.00X3, O41.00X4,
O41.00X5, O41.00X9, O41.01X0, O41.01X1, O41.01X2, O41.01X3, O41.01X4, O41.01X5, O41.01X9, O41.02X0,
O41.02X1, O41.02X2, O41.02X3, O41.02X4, O41.02X5, O41.02X9, O41.03X0, O41.03X1, O41.03X2, O41.03X3,
O41.03X4, O41.03X5, O41.03X9, O41.1010, O41.1011, O41.1012, O41.1013, O41.1014, O41.1015, O41.1019,
041.1020, 041.1021, 041.1022, 041.1023, 041.1024, 041.1025, 041.1029, 041.1030, 041.1031, 041.1032, 041.1032, 041.1032, 041.1032, 041.1032, 041.1032, 041.1032, 041.1032, 041.1033, 041.1035, 041.103
041.1033, 041.1034, 041.1035, 041.1039, 041.1090, 041.1091, 041.1092, 041.1093, 041.1094, 041.1095,
041.1099, 041.1210, 041.1211, 041.1212, 041.1213, 041.1214, 041.1215, 041.1219, 041.1220, 041.1221,
041.1222, 041.1223, 041.1224, 041.1225, 041.1229, 041.1230, 041.1231, 041.1232, 041.1233, 041.1234,
041.1235, 041.1239, 041.1290, 041.1291, 041.1292, 041.1293, 041.1294, 041.1295, 041.1299, 041.1410,
041.1411, 041.1412, 041.1413, 041.1414, 041.1415, 041.1419, 041.1420, 041.1421, 041.1422, 041.1423,
041.1424, 041.1425, 041.1429, 041.1430, 041.1431, 041.1432, 041.1433, 041.1434, 041.1435, 041.1439,

#### Pregnancy Diagnosis Code List:

The following Pregnancy Diagnosis Codes are required where indicated in the Preventive Care Services table (above), or in the Expanded Women's Preventive Health table (below):

041.1490, 041.1491, 041.1492, 041.1493, 041.1494, 041.1495, 041.1499, 041.8X10, 041.8X11, 041.8X12, 041.8X13, 041.8X14, 041.8X15, 041.8X19, 041.8X20, 041.8X21, 041.8X22, 041.8X23, 041.8X24, 041.8X25, 041.8X29, 041.8X30, 041.8X31, 041.8X32, 041.8X33, 041.8X34, 041.8X35, 041.8X39, 041.8X90, 041.8X91, 041.8X92, 041.8X93, 041.8X94, 041.8X95, 041.8X99, 041.90X0, 041.90X1, 041.90X2, 041.90X3, 041.90X4, 041.90X5, 041.90X9, 041.91X0, 041.91X1, 041.91X2, 041.91X3, 041.91X4, 041.91X5, 041.91X9, 041.92X0, 041.92X1, 041.92X2, 041.92X3, 041.92X4, 041.92X5, 041.92X9, 041.93X0, 041.93X1, 041.93X2, 041.93X3, 041.93X4, 041.93X5, 041.93X9, 042.00, 042.011, 042.012, 042.013, 042.019, 042.02, 042.10, 042.111, 042.112, 042.113, 042.119, 042.12, 042.90, 042.911, 042.912, 042.913, 042.919, 042.92, 043.011, 043.012, 043.013, 043.019, 043.021, 043.022, 043.023, 043.029, 043.101, 043.102, 043.103, 043.109, 043.111, 043.112, 043.113, 043.119, 043.121, 043.122, 043.123, 043.129, 043.191, 043.192, 043.193, 043.199, 043.211, 043.212, 043.213, 043.219, 043.221, 043.222, 043.223, 043.229, 043.231, 043.232, 043.233, 043.239, 043.811, 043.812, 043.813, 043.819, 043.891, 043.892, 043.893, 043.899, 043.90, 043.91, 043.92, 043.93, 044.00, 044.01, 044.02, 044.03, 044.10, 044.11, 044.12, 044.13, 044.20, 044.21, 044.22, 044.23, 044.30, 044.31, 044.32, 044.33, 044.40, 044.41, 044.42, 044.43, 044.50, 044.51, 044.52, 044.53, 045.001, 045.002, 045.003, 045.009, 045.011, 045.012, 045.013, 045.019, 045.021, 045.022, 045.023, 045.029, 045.091, 045.092, 045.093, 045.099, 045.8X1, 045.8X2, 045.8X3, 045.8X9, 045.90, 045.91, 045.92, 045.93, 046.001, 046.002, 046.003, 046.009, 046.011, 046.012, 046.013, 046.019, 046.021, 046.022, 046.023, 046.029, 046.091, 046.092, 046.093, 046.099, 046.8X1, 046.8X2, 046.8X3, 046.8X9, 046.90, 046.91, 046.92, 046.93, 047.00, 047.02, 047.03, 047.1, 047.9, 048.0, 048.1, 094, 098.011, 098.012, 098.013, 098.019, 098.02, 098.03, 098.111, 098.112, 098.113, 098.119, 098.12, 098.13, 098.211, 098.212, 098.213, 098.219, 098.22, 098.23, 098.311, 098.312, 098.313, 098.319, 098.32, 098.33, 098.411, 098.412, 098.413, 098.419, 098.42, 098.43, 098.511, 098.512, 098.513, 098.519, 098.52, 098.53, 098.611, 098.612, 098.613, 098.619, 098.62, 098.63, 098.711, 098.712, 098.713, 098.719, 098.72, 098.73, 098.811, 098.812, 098.813, 098.819, 098.82, 098.83, 098.911, 098.912, 098.913, 098.919, 098.92, 098.93, 099.011, 099.012, 099.013, 099.019, 099.02, 099.03, 099.111, 099.112, 099.113, 099.119, 099.12, 099.13, 099.210, 099.211, 099.212, 099.213, 099.214, 099.215, 099.280, 099.281, 099.282, 099.283, 099.284, 099.285, 099.310, 099.311, 099.312, 099.313, 099.314, 099.315, 099.320, 099.321, 099.322, 099.323, 099.324, 099.325, 099.330, 099.331, 099.332, 099.333, 099.334, 099.335, 099.340, 099.341, 099.342, 099.343, 099.344, 099.345, 099.350, 099.351, 099.352, 099.353, 099.354, 099.355, 099.411, 099.412, 099.413, 099.419, 099.42, 099.43, 099.511, 099.512, 099.513, 099.519, 099.52, 099.53, 099.611, 099.612, 099.613, 099.619, 099.62, 099.63, 099.711, 099.712, 099.713, 099.719, 099.72, 099.73, 099.810, 099.814, 099.815, 099.820, 099.824, 099.825, 099.830, 099.834, 099.835, 099.840, 099.841, 099.842, 099.843, 099.844, 099.845, 099.89, 09A.111, 09A.112, 09A.113, 09A.119, 09A.12, 09A.13, 09A.211, 09A.212, 09A.213, 09A.219, 09A.22, 09A.23, 09A.311, 09A.312, 09A.313, 09A.319, 09A.32, 09A.33, 09A.411, 09A.412, 09A.413, 09A.419, 09A.42, 09A.43, 09A.511, 09A.512, 09A.513, 09A.519, O9A.52, O9A.53, Z03.71, Z03.72, Z03.73, Z03.74, Z03.75, Z03.79, Z29.13, Z32.2, Z33.1, Z34.00, Z34.01, Z34.02, Z34.03, Z34.80, Z34.81, Z34.82, Z34.83, Z34.90, Z34.91, Z34.92, Z34.93, Z36.0, Z36.1, Z36.2, Z36.3, Z36.4, Z36.5, Z36.81, Z36.82, Z36.83, Z36.84, Z36.85, Z36.86, Z36.87, Z36.88, Z36.89, Z36.8A, Z36.9, Z3A.00, Z3A.01, Z3A.08, Z3A.09, Z3A.10, Z3A.11, Z3A.12, Z3A.13, Z3A.14, Z3A.15, Z3A.16, Z3A.17, Z3A.18, Z3A.19, Z3A.20, Z3A.21, Z3A.22, Z3A.23, Z3A.24, Z3A.25, Z3A.26, Z3A.27, Z3A.28, Z3A.29, Z3A.30, Z3A.31, Z3A.32, Z3A.33, Z3A.34, Z3A.35, Z3A.36, Z3A.37, Z3A.38, Z3A.39, Z3A.40, Z3A.41, Z3A.42, Z3A.49

#### Hepatitis C Virus Infection Screening Diagnosis Code List:

The following codes are required for the Hepatitis C Virus Infection Screening benefit. For details see the Preventive Care Services table (above).

# **ICD-10 Codes:**

A50.01,A50.02,A50.03,A50.04,A50.05,A50.06,A50.07,A50.08,A50.09,A50.1,A50.2,A50.30,A50.31,A50.32,A50.39,A 50.40,A50.41,A50.42,A50.43,A50.44,A50.45,A50.49,A50.51,A50.52,A50.53,A50.54,A50.55,A50.56,A50.57,A50.59, A50.6,A50.7,A50.9,A51.0,A51.1,A51.2,A51.31,A51.32,A51.39,A51.41,A51.42,A51.43,A51.44,A51.46,A51.49,A51.5, A51.9,A52.00,A52.01,A52.02,A52.03,A52.04,A52.05,A52.06,A52.09,A52.10,A52.11,A52.12,A52.13,A52.14,A52.15, A52.16,A52.17,A52.19,A52.2,A52.3,A52.71,A52.72,A52.73,A52.74,A52.75,A52.76,A52.77,A52.78,A52.79,A52.8,A5 2.9,A53.0,A53.9,A54.00,A54.01,A54.02,A54.03,A54.09,A54.1,A54.21,A54.22,A54.23,A54.24,A54.29,A54.30,A54.31,A54.32,A54.33,A54.39,A54.40,A54.41,A54.42,A54.43,A54.49,A54.5,A54.6,A54.81,A54.82,A54.83,A54.84,A54.85, A54.86,A54.89,A54.9,A55,A56.00,A56.01,A56.02,A56.09,A56.11,A56.19,A56.2,A56.3,A56.4,A56.8,A57,A58,A59.00,A59.01,A59.02,A59.03,A59.09,A59.8,A59.9,A60.00,A60.01,A60.02,A60.03,A60.04,A60.09,A60.1,A60.9,A63.0,A63.8,A64,A74.81,A74.89,A74.9,B07.8,B07.9,B20.,B97.35,B97.7,D65,D66,D67.,D68.0,D68.1,D68.2,D68.311,D68.312,D 68.318,D68.32,D68.4,D68.8,D68.9,F11.20,F11.21,F11.220,F11.221,F11.222,F11.229,F11.23,F11.24,F11.250,F11.251,F11.259,F11.281,F11.282,F11.288,F11.29,F12.20,F12.21,F12.220,F12.221,F12.222,F12.229,F12.250,F12.251,F

## Hepatitis C Virus Infection Screening Diagnosis Code List:

The following codes are required for the Hepatitis C Virus Infection Screening benefit. For details see the Preventive Care Services table (above).

#### ICD-10 Codes:

12,259,F12,280,F12,288,F12,29,F13,20,F13,21,F13,220,F13,221,F13,229,F13,230,F13,231,F13,232,F13,239,F13,2 4,F13.250,F13.251,F13.259,F13.26,F13.27,F13.280,F13.281,F13.282,F13.288,F13.29,F14.20,F14.21,F14.220,F14.2 21,F14.222,F14.229,F14.23,F14.24,F14.250,F14.251,F14.259,F14.280,F14.281,F14.282,F14.288,F14.29,F15.20,F1 5.21,F15.220,F15.221,F15.222,F15.229,F15.23,F15.24,F15.250,F15.251,F15.259,F15.280,F15.281,F15.282,F15.28 8,F15.29,F16.20,F16.21,F16.220,F16.221,F16.229,F16.24,F16.250,F16.251,F16.259,F16.280,F16.283,F16.288,F16. 29,F18.20,F18.21,F18.220,F18.221,F18.229,F18.24,F18.250,F18.251,F18.259,F18.27,F18.280,F18.288,F18.29,F19. 20,F19.21,F19.220,F19.221,F19.222,F19.229,F19.230,F19.231,F19.232,F19.239,F19.24,F19.250,F19.251,F19.259, F19.26,F19.27,F19.280,F19.281,F19.282,F19.288,F19.29,K50.00,K50.011,K50.012,K50.013,K50.014,K50.018,K50. 019,K50.10,K50.111,K50.112,K50.113,K50.114,K50.118,K50.119,K50.80,K50.811,K50.812,K50.813,K50.814,K50. 818,K50.819,K50.90,K50.911,K50.912,K50.913,K50.914,K50.918,K50.919,K51.20,K51.211,K51.212,K51.213,K51. 214,K51.218,K51.219,K51.30,K51.311,K51.312,K51.313,K51.314,K51.318,K51.319,K51.40,K51.411,K51.412,K51. 413,K51.414,K51.418,K51.419,K51.50,K51.511,K51.512,K51.513,K51.514,K51.518,K51.519,K51.80,K51.811,K51. 812,K51.813,K51.814,K51.818,K51.819,K51.90,K51.911,K51.912,K51.913,K51.914,K51.918,K51.919,M02.30,M02. 311,M02.312,M02.319,M02.321,M02.322,M02.329,M02.331,M02.332,M02.339,M02.341,M02.342,M02.349,M02.351 ,M02.352,M02.359,M02.361,M02.362,M02.369,M02.371,M02.372,M02.379,M02.38,M02.39,N18.3,N18.4,N18.5,N18 .6,N34.1,N49.1,N49.2,N49.3,N49.8,N49.9,N73.5,N73.9,N76.0,N76.1,N76.2,N76.3,N77.1,O35.3XX0,O35.3XX1,O35. 3XX2,O35.3XX3,O35.3XX4,O35.3XX5,O35.3XX9,O35.5XX0,O35.5XX1,O35.5XX2,O35.5XX3,O35.5XX4,O35.5XX5,O3 5.5XX9,090.4,098.011,098.012,098.013,098.019,098.02,098.03,098.111,098.112,098.113,098.119,098.12,09 8.13,098.211,098.212,098.213,098.219,098.22,098.23,098.311,098.312,098.313,098.319,098.32,098.33,099. 320,099.321,099.322,099.323,099.324,099.325,P00.2, Z00.00,Z00.01, Z04.41,Z04.42, Z05.1, Z11.3, Z11.4, Z11.59, Z11.9, Z14.01, Z14.02, Z20.2, Z20.5, Z20.6, Z20.828, Z21,Z22.4, Z41.8, Z48.21,Z48.22,Z48.24, Z48.280, Z48.288, Z48.290, Z48.298, Z49.31, Z49.32, Z51.89, Z52.000, Z52.001, Z52.008, Z52.010, Z52.011, Z52.018, Z52.090,Z52.091,Z52.098,Z52.10,Z52.11,Z52.19,Z52.20,Z52.21,Z52.29,Z52.3,Z52.4,Z52.5,Z52.6,Z52.89,Z52.9,Z 57.8, Z71.7, Z72.51, Z72.52, Z72.53, Z79.899, Z86.2, Z92.25, Z94.0, Z94.1, Z94.2, Z94.3, Z94.5, Z94.6, Z94.7, Z94.81, Z94 .82, Z94.83, Z94.84, Z94.89, Z94.9, Z95.3, Z95.4, Z99.2

## Diabetes Diagnosis Code List:

*Refer to the Preventive Care Services table (above), and the Expanded Women's Preventive Health table (below) regarding the following Diabetes Diagnosis Codes.* 

## ICD-10 Codes:

Diabetes mellitus due to underlying condition: E08.00, E08.01, E08.10, E08.11, E08.21, E08.22, E08.29, E08.311, E08.319, E08.3211, E08.3212, E08.3213, E08.3219, E08.3291, E08.3292, E08.3293, E08.3299, E08.3311, E08.3312, E08.3313, E08.3319, E08.3391, E08.3392, E08.3393, E08.3399, E08.3411, E08.3412, E08.3413, E08.3419, E08.3491, E08.3492, E08.3493, E08.3499, E08.3511, E08.3512, E08.3513, E08.3519, E08.3521, E08.3522, E08.3523, E08.3529, E08.3531, E08.3532, E08.3533, E08.3539, E08.3541, E08.3542, E08.3543, E08.3549, E08.3551, E08.3552, E08.3553, E08.3559, E08.3591, E08.3592, E08.3593, E08.3599, E08.3599, E08.36, E08.37X1, E08.37X2, E08.37X3, E08.37X9, E08.39, E08.40, E08.41, E08.42, E08.43, E08.44, E08.49, E08.51, E08.52, E08.610, E08.618, E08.620, E08.621, E08.622, E08.628, E08.630, E08.638, E08.641, E08.649, E08.65, E08.69, E08.8, E08.9

Drug or chemical induced diabetes mellitus: E09.00, E09.01, E09.10, E09.11, E09.21, E09.22, E09.29, E09.311, E09.319, , E09.3211, E09.3212, E09.3213, E09.3219, , E09.3291, E09.3292, E09.3293, E09.3299, E09.3311, E09.3312, E09.3313, E09.3319, E09.3391, E09.3392, E09.3393, E09.3399, E09.3411, E09.3412, E09.3413, E09.3419, E09.3491, E09.3492, E09.3493, E09.3499, E09.3511, E09.3512, E09.3513, E09.3519, E09.3521, E09.3522, E09.3523, E09.3529, E09.3531, E09.3532, E09.3533, E09.3539, E09.3541, E09.3542, E09.3543, , E09.3549, E09.3551, E09.3552, E09.3553, E09.3559, E09.3591, E09.3593, E09.3593, E09.3599, E09.36, E09.37X1, E09.37X2, E09.37X3, E09.37X9, E09.39, E09.40, E09.41, E09.42, E09.43, E09.44, E09.49, E09.51, E09.52, E09.59, E09.610, E09.618, E09.620, E09.621, E09.622, E09.628, E09.630, E09.638, E09.641, E09.649, E09.65, E09.69, E09.8, E09.9

Type 1 diabetes mellitus: E10.10, E10.11, E10.21, E10.22, E10.29, E10.311, E10.319, E10.3211, E10.3212, E10.3213, E10.3219, E10.3291, E10.3292, E10.3293, E10.3299, E10.3311, E10.3312, E10.3313, E10.3319, E10.3391, E10.3392, E10.3393, E10.3399, E10.3411, E10.3412, E10.3413, E10.3419, E10.3491, E10.3492, E10.3493, E10.3499, E10.3511, E10.3512, E10.3513, E10.3519, E10.3521, E10.3522, E10.3523, E10.3529, E10.3531, E10.3532, E10.3533, E10.3539, E10.3541, E10.3542, E10.3543, E10.3549, E10.3551, E10.3552, E1

#### Diabetes Diagnosis Code List:

Refer to the Preventive Care Services table (above), and the Expanded Women's Preventive Health table (below) regarding the following Diabetes Diagnosis Codes.

E10.3553, E10.3559, E10.3591, E10.3592, E10.3593, E10.3599, E10.36, E10.37X1, E10.37X2, E10.37X3, E10.37X9, E10.39, E10.40, E10.41, E10.42, E10.43, E10.44, E10.49, E10.51, E10.52, E10.59, E10.610, E10.618, E10.620, E10.621, E10.622, E10.628, E10.630, E10.638, E10.641, E10.649, E10.65, E10.69, E10.8, E10.9

Type 2 diabetes mellitus: E11.00, E11.01, E11.10, E11.11, E11.21, E11.22, E11.29, E11.311, E11.319, E11.3211, E11.3212, E11.3213, E11.3219, E11.3291, E11.3292, E11.3293, E11.3299, E11.3311, E11.3312, E11.3313, E11.3319, E11.3392, E11.3393, E11.3399, E11.3411, E11.3412, E11.3413, E11.3419, E11.3491, E11.3492, E11.3493, E11.3499, E11.3511, E11.3512, E11.3513, E11.3519, E11.3521, E11.3522, E11.3523, E11.3529, E11.3531, E11.3532, E11.3533, E11.3539, E11.3541, E11.3542, E11.3543, E11.3549, E11.3551, E11.3552, E11.3553, E11.3559, E11.3591, E11.3592, E11.3593, E11.3599, E11.36, E11.37X1, E11.37X2, E11.37X3, E11.37X9, E11.39, E11.40, E11.41, E11.42, E11.43, E11.44, E11.49, E11.51, E11.52, E11.59, E11.610, E11.618, E11.620, E11.621, E11.622, E11.628, E11.630, E11.638, E11.641, E11.649, E11.65, E11.69, E11.8, E11.9

Other specified diabetes mellitus: E13.00, E13.01, E13.10, E13.11, E13.21, E13.22, E13.29, E13.311, E13.319, E13.3211, E13.3212, E13.3213, E13.3219, E13.3291, E13.3292, E13.3293, E13.3299, E13.3311, E13.3312, E13.3313, E13.3319, E13.3391, E13.3392, E13.3393, E13.3399, E13.3411, E13.3412, E13.3413, E13.3419, E13.3491, E13.3492, E13.3493, E13.3499, E13.3511, E13.3512, E13.3513, E13.3519, E13.3521, E13.3522, E13.3523, E13.3529, E13.3531, E13.3532, E13.3539, E13.3541, E13.3542, E13.3543, E13.3549, E13.3551, E13.3552, E13.3553, E13.3559, E13.3591, E13.3592, E13.3593, E13.3599, E13.3599, E13.36, E13.37X1, E13.37X2, E13.37X3, E13.37X9, E13.39, E13.40, E13.41, E13.42, E13.43, E13.44, E13.49, E13.51, E13.52, E13.59, E13.610, E13.618, E13.620, E13.621, E13.622, E13.628, E13.630, E13.638, E13.641, E13.649, E13.65, E13.69, E13.8, E13.9

#### Atherosclerosis Diagnosis Code List:

Refer to the Preventive Care Services table (above), regarding the following Atherosclerosis Diagnosis Codes which are allowed for:

- Cholesterol Screening (Lipid Disorders Screening)
- Behavioral Counseling in Primary Care to Promote a Healthful Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors

## **ICD-10 Codes:**

170.0, 170.1, 170.201, 170.202, 170.203, 170.208, 170.209, 170.211, 170.212, 170.213, 170.218, 170.219, 170.221, 170.222, 170.223, 170.228, 170.229, 170.231, 170.232, 170.233, 170.234, 170.235, 170.238, 170.239, 170.241, 170.242, 170.243, 170.244, 170.245, 170.248, 170.249, 170.25, 170.261, 170.262, 170.263, 170.268, 170.269, 170.291, 170.292, 170.293, 170.298, 170.299, 170.301, 170.302, 170.303, 170.308, 170.309, 170.311, 170.312, 170.313, 170.318, 170.319, 170.321, 170.322, 170.323, 170.328, 170.329, 170.331, 170.332, 170.333, 170.334, 170.335, 170.338, 170.339, 170.341, 170.342, 170.343, 170.344, 170.345, 170.348, 170.349, 170.35, 170.361, 170.362, 170.363, 170.368, 170.369, 170.391, 170.392, 170.393, 170.398, 170.399, 170.401, 170.402, 170.403, 170.408, 170.409, 170.411, 170.412, 170.413, 170.418, 170.419, 170.421, 170.422, 170.423, 170.428, 170.429, 170.431, 170.432, 170.433, 170.434, 170.435, 170.438, 170.439, 170.441, 170.442, 170.443, 170.444, 170.445, 170.448, 170.449, 170.45, 170.461, 170.462, 170.463, 170.468, 170.469, 170.491, 170.492, 170.493, 170.498, 170.499, 170.501, 170.502, 170.503, 170.508, 170.509, 170.511, 170.512, 170.513, 170.518, 170.519, 170.521, 170.522, 170.523, 170.528, 170.529, 170.531, 170.532, 170.533, 170.534, 170.535, 170.538, 170.539, 170.541, 170.542, 170.543, 170.544, 170.545, 170.548, 170.549, 170.55, 170.561, 170.562, 170.563, 170.568, 170.569, 170.591, 170.592, 170.593, 170.598, 170.599, 170.601, 170.602, 170.603, 170.608, 170.609, 170.611, 170.612, 170.613, 170.618, 170.619, 170.621, 170.622, 170.623, 170.628, 170.629, 170.631, 170.632, 170.633, 170.634, 170.635, 170.638, 170.639, 170.641, 170.642, 170.643, 170.644, 170.645, 170.648, 170.649, 170.65, 170.661, 170.662, 170.663, 170.668, 170.669, 170.691, 170.692, 170.693, 170.698, 170.699, 170.701, 170.702, 170.703, 170.708, 170.709, 170.711, 170.712, 170.713, 170.718, 170.719, 170.721, 170.722, 170.723, 170.728, 170.729, 170.731, 170.732, 170.733, 170.734, 170.735, 170.738, 170.739, 170.741, 170.742, 170.743, 170.744, 170.745, 170.748, 170.749, 170.75, 170.761, 170.762, 170.763, 170.768, 170.769, 170.791, 170.792, 170.793, 170.798, 170.799, 170.8, 170.90, 170.91

# These are the requirements of Health and Human Services for plan years that begin on or after 8/1/12.

For additional services covered for women, see the Preventive Care Services table above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service:	Code(s):	Preventive Benefit Instructions:
Well-Woman Visits HHS Requirement: Well-woman preventive care visit annually for adult women to obtain the recommended preventive services that are age and developmentally appropriate, including preconception and prenatal care.	<ul> <li>Procedure Code(s): <u>Well-woman visits:</u> <ul> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> </ul> </li> <li><u>Prenatal Office Visits:</u> Evaluation and Management (Office Visits): <ul> <li>99201, 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0463</li> </ul> </li> </ul>	<ul> <li><u>Well-woman visits:</u></li> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> <li><u>Prenatal Office Visits:</u></li> <li>Payable as preventive when billed with a Pregnancy Diagnosis Code (see Pregnancy Diagnosis Code list above).</li> </ul>
Also see "Wellness Examinations" section in the Preventive Care Services table above.	<ul> <li>Physician prenatal education, group setting:</li> <li>99078</li> <li><u>Prenatal Care Visits:</u></li> <li>59425, 59426</li> <li><u>Global Obstetrical Codes:</u></li> <li>59400, 59510, 59610, 59618</li> <li><b>Diagnosis Code(s):</b></li> <li>See Pregnancy Diagnosis Code list above.</li> </ul>	<ul> <li>Prenatal Care Visits:</li> <li>Pregnancy Diagnosis Codes are not required.</li> <li>Global Obstetrical Codes:</li> <li>The routine, low-risk, prenatal visits portion of the code is covered as preventive.</li> <li>Pregnancy Diagnosis Codes are not required.</li> </ul>
Screening for Gestational Diabetes HHS Requirement: Women who are 24 to 28 weeks pregnant, and at the first prenatal visit for those who are at high risk of development of gestational diabetes. Also see Diabetes Screening and the Gestational Diabetes Mellitus Screening sections in the Preventive Care Services table above.	<ul> <li>Procedure Code(s): <ul> <li>82947, 82948, 82950, 82951, 82952, 83036</li> <li>36415, 36416</li> </ul> </li> <li>Diagnosis Code(s): <ul> <li>Pregnancy Diagnosis Code. (See Pregnancy Diagnosis Code list above.)</li> </ul> </li> </ul>	<ul> <li>Payable with Pregnancy Diagnosis Code (regardless of gestational week)</li> <li><u>Criteria for 36415 and 36416</u>: Payable when billed with <b>ALL</b> of the following:         <ul> <li>With one of the Diabetes Screening Procedure codes listed in this row</li> <li><u>AND</u></li> <li>With a Pregnancy Diagnosis Code</li> </ul> </li> <li>NOTE: If a Diabetes Diagnosis Code is present in any position, the preventive benefit will <u>not</u> be applied. See Diabetes Diagnosis Codes table above.</li> </ul>
Human Papillomavirus DNA Testing (HPV) HHS Requirement: High-risk human papillomavirus DNA testing in women with normal cytology (pap smear) results, every 3 years for women who are 30 or older.	<pre>Procedure Code(s):</pre>	<ul> <li>Age 30 and up.</li> <li>Payable as a preventive screening with one of the Diagnosis Codes listed in this row.</li> </ul>

These are the requirements of Health and Human Services for plan years that begin on or after 8/1/12.

For additional services covered for women, see the Preventive Care Services table above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service:	Code(s):	Preventive Benefit Instructions:
Counseling for Sexually Transmitted Infections HHS Requirement: Counseling on sexually transmitted infections for all sexually active women.	<ul> <li>See the Wellness</li> <li>Examinations section of the Preventive Care Services table above.</li> </ul>	<ul> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> </ul>
Counseling and Screening for Human Immune-deficiency Virus HHS Requirement: Counseling and screening for human immune-deficiency virus infection for all sexually active women.	<ul> <li>Counseling:</li> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> <li>Screening Tests:</li> <li>See the HIV - Human Immunodeficiency Virus - Screening for Adolescents and Adults section of the Preventive Services table above.</li> </ul>	<ul> <li>Counseling:</li> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> <li>Screening Tests:</li> <li>See the HIV – Human Immunodeficiency Virus –Screening for Adolescents and Adults section of the Preventive Services table above.</li> </ul>
Contraceptive Methods (Including Sterilizations) HHS Requirement: For women, all Food and Drug Administration-approved contraceptive methods, sterilization procedures, and patient education and counseling (as prescribed).	Code Group 1: <u>Sterilizations:</u> • Tubal Ligation, oviduct occlusion: 58565, 58600, 58605, 58611, 58615, 58670, 58671, A4264 (See Code Group 4 below for tubal ligation followup.)	<ul> <li>Code Group 1 :</li> <li>Does not have diagnosis code requirements for preventive benefits to apply.</li> <li>Code Group 2:</li> <li>Preventive when billed with one of the Code Group 2 Diagnosis Codes, listed in this row.</li> </ul>
For patient education and counseling: • See the Wellness Examinations section of the Preventive Care Services table above. <u>NOTES</u> : Certain employers may qualify for an exemption from covering contraceptive methods and sterilizations on account of religious objections.	<ul> <li><u>Contraceptive Methods:</u></li> <li>Diaphragm or cervical cap: 57170, A4261, A4266</li> <li>IUD (copper): J7300</li> <li>IUD (Skyla<sup>®</sup>): J7301</li> <li>IUD (Liletta<sup>®</sup>): J7297</li> <li>IUD (Kyleena<sup>®</sup>): Q9984</li> <li>See Code Group 2 for <u>additional</u> <u>IUD</u> codes.</li> </ul> Code Group 2: Contraceptive Methods: <ul> <li>Implantable devices:</li> </ul>	<ul> <li>Code Group 3:</li> <li>Preventive when billed with the Code Group 3 Diagnosis Code listed in this row.</li> <li>Code Group 4:</li> <li>Preventive when billed with the Code Group 4 Diagnosis Code listed in this row.</li> </ul>
Refer to the Outpatient Prescription Drug Rider, or SPD for self funded plans, for specific prescription drug product coverage and exclusion terms, and myuhc.com for information regarding coverage for contraceptive drugs.	<ul> <li>11981 (implant insertion)</li> <li>11982 (implant removal)</li> <li>11983 (removal with reinsertion)</li> </ul>	<b>Code Group 5:</b> • Preventive when billed with the Code Group 5 Diagnosis code listed in this row.

Preventive Care Services

Page 43 of 50 Effective 10/01/2017

UnitedHealthcare Commercial Coverage Determination Guideline Effective Proprietary Information of UnitedHealthcare. Copyright 2017 United HealthCare Services, Inc.

## These are the requirements of Health and Human Services for plan years that begin on or after 8/1/12.

For additional services covered for women, see the Preventive Care Services table above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service:	Code(s):	Preventive Benefit Instructions:
	<u>additional IUD</u> codes.	
	<ul> <li>Injections:         <ul> <li>J1050 (injection)</li> <li>96372 (administration)</li> </ul> </li> </ul>	
	Code Group 2 Diagnosis Code(s) These are required for Code Group 2:	
	Contraceptive Management: <u>ICD-10:</u> Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9	
	<b>Code Group 3:</b> Anesthesia for Sterilization: • 00851, 00940, 00942, 00950, 00952, 01960, 01961, 01965, 01966, 01967, 01968	
	Code Group 3 Diagnosis Code: <i>Sterilization:</i> • <u>ICD-10:</u> Z30.2	
	<b>Code Group 4:</b> Tubal ligation followup hysterosalpingogram:	
	<i>Catheterization and introduction of saline or contrast material:</i> 58340	
	Hysterosalpingography: 74740 Contrast material: Q9967	
	Code Group 4 Diagnosis Code: <i>Tubal ligation status:</i> • <u>ICD-10:</u> Z98.51	
	Code Group 5: IUD Followup Visit: • 99211, 99212	
	Code Group 5 Diagnosis Code: • <u>ICD-10:</u> Z30.431	

These are the requirements of Health and Human Services for plan years that begin on or after 8/1/12.

For additional services covered for women, see the Preventive Care Services table above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service:	Code(s):	Preventive Benefit Instructions:
Breastfeeding Support, Supplies, and Counseling HHS Requirement: Breastfeeding support, supplies, and counseling: Comprehensive lactation support and counseling, from a trained provider, during pregnancy and/or in the postpartum period, and costs for renting breastfeeding equipment, in conjunction with each birth.	Support and Counseling: Procedure Code(s): • S9443 • 99241, 99242, 99243, 99244, 99245 • 99341, 99342, 99343, 99344, 99345 • 99347, 99348, 99349, 99350 Also see the codes in the Wellness Examinations section of the Preventive Care Services table above. Diagnosis Code(s): • ICD-10: Z39.1 Breast Pump Equipment & Supplies: Procedure Code(s): Personal Use Electric: • E0603 Breast Pump Supplies: • A4281, A4282, A4283, A4284, A4285, A4286 Diagnosis Code(s): • Pregnancy Diagnosis Code (see Pregnancy Diagnosis Code list above), OR • ICD-10: Z39.1	<ul> <li>Support and Counseling:         <ul> <li>The Diagnosis Code listed in this row is required for 99241 – 99245, 99341 – 99345, and 99347 – 99350</li> <li>The Diagnosis Code listed in this row is not required for S9443</li> </ul> </li> <li>Breast Pump Equipment &amp; Supplies:         <ul> <li>E0603 is limited to one purchase per birth.</li> <li>E0603, and A4281 – A4286 are payable as preventive with at least one of the diagnosis codes listed in this row.</li> </ul> </li> </ul>
Screening and Counseling for Interpersonal and Domestic Violence HHS Requirement: Screening and counseling for interpersonal and domestic violence.	<ul> <li>Procedure Code(s):</li> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> <li>Diagnosis Code(s):</li> <li>See the Wellness Examinations section of the Preventive Care Services table above.</li> </ul>	• n/a

#### Revenue Code

See Mammography and Immunizations sections above for the applicable revenue codes.

#### REFERENCES

ACIP Abbreviations for Vaccines (including Trade Names): <u>http://www.cdc.gov/vaccines/acip/committee/guidance/vac-abbrev.html</u>. Accessed June 7, 2017.

ACIP Recommendations: <u>http://www.cdc.gov/vaccines/hcp/acip-recs/vacc-specific/index.html</u>. Accessed June 7, 2017.

American Academy of Family Physicians (AAFP) Summary of Recommendations for Clinical Preventive Services: <u>http://www.aafp.org/dam/AAFP/documents/patient\_care/clinical\_recommendations/cps-recommendations.pdf</u>. Accessed June 7, 2017.

American Academy of Pediatrics / Bright Futures / Recommendations for Pediatric Preventive Healthcare. (For ages 0 – 21): <u>http://www.aap.org/en-us/professional-resources/practice-</u>support/Periodicity/Periodicity%20Schedule FINAL.pdf. Accessed June 7, 2017.

American Academy of Pediatrics: <u>http://www.aap.org/</u>. Accessed June 7, 2017.

Centers for Disease Control and Prevention / Immunization Schedules: <u>http://www.cdc.gov/vaccines/schedules/index.html</u>. Accessed June 7, 2017.

Department of Labor: <u>http://www.dol.gov/ebsa/consumer\_info\_health.html</u>. Accessed June 7, 2017.

Grade Definitions for USPSTF Recommendations <u>http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions</u>. Accessed June 7, 2017.

July 19, 2010 IRS Interim Rules: http://www.irs.gov/irb/2010-29 IRB/index.html. Accessed June 7, 2017.

Published Recommendations, U.S. Preventive Services Task Force: <u>http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations</u>. Accessed June 7, 2017.

U.S. Food and Drug Administration (FDA), Vaccines Licensed for Use in the United States: <u>http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM093833</u>. Accessed June 7, 2017.

Women's Preventive Services: Required Health Plan Coverage Guidelines: <u>http://www.hrsa.gov/womensguidelines/</u>. Accessed June 7, 2017.

# GUIDELINE HISTORY/REVISION INFORMATION

Date	Action/Description
10/01/2017	<ul> <li>Updated coverage rationale/indications for coverage:</li> <li>Pediatrics         <ul> <li>Clarified list of services covered under the preventive care benefit for children (of the appropriate age); replaced "counseling for fluoride for prevention of dental cavities" with "application of fluoride by a primary care provider for prevention of dental cavities"</li> </ul> </li> <li>Added language to indicate the definitions listed in the policy may not apply to all plans; refer to the member specific benefit plan document for applicable definitions</li> <li>Revised list of applicable procedure and diagnosis codes for:</li> <li>Preventive Care Services</li> <li>Chlamydia Infection Screening</li> <li>Added notation to indicate Bright Futures recommends sexually transmitted infection screening if risk assessment is positive between age 11–21 years</li> <li>Gonorrhea Screening</li> <li>Added notation to indicate Bright Futures recommends sexually transmitted infection screening if risk assessment is positive between age 11–21 years</li> <li>HUV – Human Immunodeficiency Virus – Screening for Adolescents and Adults</li> <li>Added notation to indicate Bright Futures recommends HIV screening lab work:         <ul> <li>Once between age 15–18 years</li> <li>Anytime between age 11–14 years and 19–21 years if risk assessment is positive</li> </ul> </li> <li>Cholesterol Screening (Lipid Disorders Screening)</li> <li>Added instruction to refer to the Dyslipidemia Screening (Bright Futures) section of the policy for recommendations for children Newborn Screenings</li> <li>Added instruction to refer to the Hearing Tests (Bright Futures) section of the policy for information on hearing screening for newborns</li> <li>Screening and Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse</li> <li>Removed/replaced notation indicating the Bright Futures Periodicity Schedule recommends</li></ul>

#### Action/Description

• Annual alcohol screening (HCPCS code G0442)

Brief counseling for alcohol (HCPCS code G0443)

- Depression in Children and Adolescents (Screening)
- Replaced notation indicating "the Bright Futures Periodicity Schedule recommends depression screening begin at age 12" with "the Bright Futures Periodicity Schedule recommends depression screening begin at age 12 through 21 years"

Primary Care Interventions to Prevent Tobacco Use in Children and Adolescents

- Added notation to indicate Bright Futures (April 2017) recommends tobacco use assessments from age 11–21 years
- Reformatted list of applicable procedure codes; added code classification/sub-heading for "smoking and tobacco use cessation counseling visit" (CPT codes 99401, 99402, 99403, 99404, 99406, and 99407)

#### Preeclampsia Screening (new to policy)

- Added language to indicate:
  - The USPSTF recommends screening for preeclampsia in pregnant women with blood pressure measurements throughout pregnancy (USPSTF Rating: B)
  - Preeclampsia screening by blood pressure measurement is included in the code for a wellness examination visit; see the *Wellness Examinations* section of the policy for applicable procedure codes and preventive benefit instructions

## **Bright Futures**

Hearing Tests

- Added language to indicate Bright Futures (April 2017) recommends:
  - Hearing test:
    - At age newborn, between 3-5 days to 2 months, 4 years, 5 years, 6 years, 8 years, 10 years, once between 11–14 years, once between 15–17 years, and once between 18–21 years
    - If risk assessment is positive
  - Risk assessment:
    - At age 4 months, 6 months, 9 months, 12 months, 15 months, 18 months, 24 months, 30 months, 3 years, 7 years, and 9 years
- Updated and reformatted list of applicable procedure codes:
  - Added code classification/sub-heading for "hearing tests" (CPT codes 92551, 92552, and 92553)
  - Added notation to indicate a risk assessment is included in the code for a wellness examination visit; see the *Wellness Examinations* section of the policy for applicable procedure codes

Dyslipidemia Screening

- Added language to indicate Bright Futures (April 2017) recommends:
- Risk assessment:
  - At age 24 months, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, and 16 years
  - Lab work:

0

- If risk assessment is positive; or
- Once between age 9–11 years and once between age 17–21 years
- Updated and reformatted list of applicable procedure codes:
  - Modified code classification/sub-heading; replaced "dyslipidemia screening" with "dyslipidemia screening *lab work*"
  - Added notation to indicate a risk assessment is included in the code for a wellness examination visit; see the *Wellness Examinations* section of the policy for applicable procedure codes

Tobacco, Alcohol or Drug Use Assessment (new to policy)

- Added language to indicate Bright Futures (April 2017) recommends tobacco, alcohol or drug use assessment from age 11 years-21 years
- Added instruction to see the following sections of the policy for applicable procedure codes and preventive benefit instructions:
  - Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents and Screening

Action/Description
<ul> <li>Behavioral Counseling Interventions in Primary Care to Reduce Alcohol Misuse</li> </ul>
Psychosocial/Behavioral Assessment Bright Futures (new to policy) <ul> <li>Added language to indicate:</li> </ul>
<ul> <li>Bright Futures (April 2017) recommends psychosocial/behavioral assessment at each of the recommended visits between age newborn- 21 years</li> </ul>
<ul> <li>An assessment is included in the procedure code for a wellness examination visit</li> </ul>
<ul> <li>Added instruction to see the Wellness Examinations section of the policy for applicable procedure codes and preventive benefit instructions</li> </ul>
Depression Screening (new to policy)
<ul> <li>Added language to indicate Bright Futures (April 2017) recommends depression screening at each of the recommended visits between age 12 years-21 years</li> </ul>
<ul> <li>Added instruction to see the <i>Depression in Children and Adolescents</i> (<i>Screening</i>) section of the policy for applicable procedure codes and preventive benefit instructions</li> </ul>
Sexually Transmitted Infections (STI) (new to policy)
<ul> <li>Added language to indicate Bright Futures (April 2017) recommends:</li> </ul>

- Risk assessment:
  - At each of the recommended visits between age 11–21 years
- Lab work:

Date

- If risk assessment is positive
- Added instruction to see the following sections of the policy for applicable procedure codes and preventive benefit instructions:
  - Chlamydia Infection Screening
  - Gonorrhea Screening

HIV Screening (new to policy)

• Added language to indicate Bright Futures (April 2017) recommends:

- Risk assessment:
  - At age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years, and 21 years
- Lab work:
  - Once between age 15–18 years
  - Anytime between age 11–14 years and 19–21 years when risk assessment is positive
- Added instruction to see the *HIV Human Immunodeficiency Virus Screening for Adolescents and Adults* section of the policy for applicable procedure codes and preventive benefit instructions

## **Pregnancy Diagnosis Code List**

- Updated list of applicable ICD-10 diagnosis codes to reflect annual code edits:
  - Added 000.101, 000.102, 000.109, 000.111, 000.112, 000.119, 000.201, 000.202, 000.209, 000.211, 000.212, 000.219, 036.8310, 036.8311, 036.8312, 036.8313, 036.8314, 036.8315, 036.8319, 036.8320, 036.8321, 036.8322, 036.8323, 036.8324, 036.8325, 036.8329, 036.8330, 036.8331, 036.8332, 036.8333, 036.8334, 036.8335, 036.8339, 036.8390, 036.8391, 036.8392, 036.8393, 036.8394, 036.8395, 036.8399, Z36.0, Z36.1, Z36.2, Z36.3, Z36.4, Z36.5, Z36.81, Z36.82, Z36.83, Z36.84, Z36.85, Z36.86, Z36.87, Z36.88, Z36.89, Z36.8A, and Z36.9
  - Removed 000.10, 000.11, 000.20, 000.21, and Z36

#### **Diabetes Diagnosis Code List**

- Updated list of applicable ICD-10 diagnosis codes to reflect annual code edits; added E11.10 and E11.11
- Archived previous policy version CDG.016.16

# Appendix A – USPSTF Grade Definitions

# Grade Definitions for USPSTF Recommendations <u>After</u> July 2012:

http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions

Grade	Definition	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
В	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
С	The USPSTF recommends selectively offering or providing this service to individual patients based on professional judgment and patient preferences. There is at least moderate certainty that the net benefit is small.	Offer or provide this service for selected patients depending on individual circumstances.
D	The USPSTF recommends against the service. There is moderate or high certainty that the service has no net benefit or that the harms outweigh the benefits.	Discourage the use of this service.
I	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

# Grade Definitions for USPSTF Recommendations Dated After May 2007:

http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions

The U.S. Preventive Services Task Force (USPSTF) has updated its definitions of the grades it assigns to recommendations and now includes "suggestions for practice" associated with each grade. The USPSTF has also defined levels of certainty regarding net benefit. These definitions apply to USPSTF recommendations voted on after May 2007.

Grade	Definition	Suggestions for Practice
A	The USPSTF recommends the service. There is high certainty that the net benefit is substantial.	Offer or provide this service.
В	The USPSTF recommends the service. There is high certainty that the net benefit is moderate or there is moderate certainty that the net benefit is moderate to substantial.	Offer or provide this service.
С	Note: The following statement is undergoing revision. Clinicians may provide this service to selected patients depending on individual circumstances. However, for most individuals without signs or symptoms there is likely to be only a small benefit from this service.	Offer or provide this service only if other considerations support the offering or providing the service in an individual patient.
D	Offer or provide this service only if other considerations support the offering or providing the service in an individual patient.	Discourage the use of this service.
I	The USPSTF concludes that the current evidence is insufficient to assess the balance of benefits and harms of the service. Evidence is lacking, of poor quality, or conflicting, and the balance of benefits and harms cannot be determined.	Read the clinical considerations section of USPSTF Recommendation Statement. If the service is offered, patients should understand the uncertainty about the balance of benefits and harms.

# **Grade Definitions Prior to May 2007**

The definitions below (of USPSTF grades and quality of evidence ratings) were in use prior to the update and apply to recommendations voted on by the USPSTF prior to May 2007.

**A - Strongly Recommended:** The USPSTF strongly recommends that clinicians provide [the service] to eligible patients. *The USPSTF found good evidence that [the service] improves important health outcomes and concludes that benefits substantially outweigh harms*.

**B** - **Recommended:** The USPSTF recommends that clinicians provide [the service] to eligible patients. *The USPSTF found at least fair evidence that [the service] improves important health outcomes and concludes that benefits outweigh harms*.

**C** - **No Recommendation:** The USPSTF makes no recommendation for or against routine provision of [the service]. *The USPSTF found at least fair evidence that [the service] can improve health outcomes but concludes that the balance of benefits and harms is too close to justify a general recommendation.* 

**D** - **Not Recommended:** The USPSTF recommends against routinely providing [the service] to asymptomatic patients. *The USPSTF found at least fair evidence that [the service] is ineffective or that harms outweigh benefits.* 

**I** - **Insufficient Evidence to Make a Recommendation:** The USPSTF concludes that the evidence is insufficient to recommend for or against routinely providing [the service]. *Evidence that the [service] is effective is lacking, of poor quality, or conflicting and the balance of benefits and harms cannot be determined*.